

Question Paper Preview

Notations :

- 1.Options shown in green color and with ✓ icon are correct.
- 2.Options shown in red color and with ✗ icon are incorrect.

Question Paper Name:	Civil Assistant Surgeon in ENT 5th Dec 2017 Shift 1
Subject Name:	Civil Assistant Surgeon in ENT
Creation Date:	2017-12-05 14:04:25
Duration:	150
Total Marks:	150
Display Marks:	No
Calculator:	Scientific
Magnifying Glass Required?:	No
Ruler Required?:	No
Eraser Required?:	No
Scratch Pad Required?:	No
Rough Sketch/Notepad Required?:	No
Protractor Required?:	No

Civil Assistant Surgeon in ENT

Group Number :	1
Group Id :	57996565
Group Maximum Duration :	0
Group Minimum Duration :	150
Revisit allowed for view? :	No
Revisit allowed for edit? :	No
Break time:	0
Group Marks:	150

Civil Assistant Surgeon in ENT

Section Id :	57996565
Section Number :	1
Section type :	Online
Mandatory or Optional:	Mandatory
Number of Questions:	150
Number of Questions to be attempted:	150
Section Marks:	150
Display Number Panel:	Yes
Group All Questions:	No

Sub-Section Number:	1
Sub-Section Id:	579965103
Question Shuffling Allowed :	Yes

Question Number : 1 Question Id : 5799659653 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Distance of carina from external nares is

Options :

- ✓ 11.8 inches
- ✗ 9.8 inches
- ✗ 7.9 inches
- ✗ 5.9 inches

Question Number : 2 Question Id : 5799659654 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Early acute suppurative otitis media is manifested by:

Options :

- ✗ Retracted tympanic membrane
- ✗ Aural fullness
- ✗ Deafness
- ✓ All the given options are correct

Question Number : 3 Question Id : 5799659655 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False statement about autoimmune ear disease is

Options :

- ✗ Histopathology from suspected patients shows fibrosis and/or bone deposition in labyrinth
- ✓ Lymphocytic migration assays using inner ear tissues as target have been very promising.
- ✗ Suspected patients show reactivity against a 68 kD antigen.
- ✗ It is associated with reactivity against 45 to 50 kD, 30 kD and 20 kD antigens.

Question Number : 4 Question Id : 5799659656 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Parkinson's disease is characterized by

Options :

- ✗ Hypophonia is a hallmark in late parkinson's disease
- ✓ Dysarthria becomes severe with progression of disease
- ✗ Patient can voluntarily produce phonation even in late stages
- ✗ Percutaneous collagen improves overall speech in patients with pre-existing dysarthria.

Question Number : 5 Question Id : 5799659657 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following organisms are common sources of infection in acute Bacterial sialadenitis?

Options :

- ✗ Gram-positive cocci
- ✗ Aerobic gram-negative rods
- ✗ Anaerobic gram-negative rods
- ✓ All the given options are correct

Question Number : 6 Question Id : 5799659658 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The middle ear magnifies the sound:

Options :

1. ✘ 20 times
2. ✔ 22 times
3. ✘ 200 times
4. ✘ 21 times

Question Number : 7 Question Id : 5799659659 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False statement about allergic rhinitis is

Options :

1. ✘ About 75% of patients with seasonal allergens have symptoms attributable to ragweed.
2. ✘ Dust mite faeces, an allergen remains airborne for short periods.
3. ✘ Dust mite faeces cannot be effectively removed by air filtration systems.
4. ✔ Aspergillus and Penicillium molds are principally responsible for symptoms attributable to outdoor exposure.

Question Number : 8 Question Id : 5799659660 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a manifestation of hypercalcemia?

Options :

1. ✘ Confusion
2. ✘ Polyuria
3. ✘ Nephrolithiasis
4. ✔ Tetany

Question Number : 9 Question Id : 5799659661 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False statement about tympanometry is

Options :

1. ✘ Used as screening and diagnostic tool of peripheral and central auditory disorders
2. ✘ Classification was first described by Liden.
3. ✘ Type D pattern is often seen with scarred eardrums or with normal hypermobile eardrums.
4. ✔ The neural network of the acoustic reflex is located in the upper brainstem.

Question Number : 10 Question Id : 5799659662 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The most commonly affected site for laryngeal sarcoid is:

Options :

1. ✘ Subglottis
2. ✘ Glottis
3. ✔ Supraglottis
4. ✘ Anterior Commissure

Question Number : 11 Question Id : 5799659663 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The earliest manifestation of cavernous sinus thrombosis is:

Options :

- ✘ Fever.
- ✘ Ptosis.
- ✘ Proptosis.
- ✔ Ophthalmoplegia

Question Number : 12 Question Id : 5799659664 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A pregnant (20 weeks, primipara) woman presents to your clinic complaining of severe nasal congestion for the past month, which has made it difficult for her to sleep. On examination she has a straight septum, bilateral boggy inferior turbinates, and no additional nasal abnormalities on endoscopy. What is the best initial recommendation?

Options :

- ✔ Trial of saline irrigations.
- ✘ Loratadine (Claritin) and a medrol dose pack
- ✘ Loratadine (Claritin) and steroid nasal sprays.
- ✘ Do nothing, as there are no safe medications for the developing fetus

Question Number : 13 Question Id : 5799659665 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Antrum of Highmore is another name for:

Options :

- ✔ maxillary
- ✘ frontal
- ✘ Sphenoid
- ✘ Ethmoid

Question Number : 14 Question Id : 5799659666 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False statement regarding retropharyngeal abscess is:

Options :

- ✘ It lies lateral to midline
- ✘ Causes difficulty in swallowing and speech.
- ✘ Can always be palpated by finger at posterior pharyngeal wall.
- ✔ It is present beneath the vertebral fascia.

Question Number : 15 Question Id : 5799659667 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A 10 year old boy developed hoarseness of voice following an attack of diphtheria. On examination, his Rt vocal cord was paralysed. The treatment of choice for paralysed vocal cord will be:

Options :

- ✘ Gel foam injection of right vocal cord

- ✘ Fat injection of right vocal cord
- ✘ Thyroplasty type—I
- ✔ Wait for spontaneous recovery of vocal cord

Question Number : 16 Question Id : 5799659668 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A 55 year old female presents with tinnitus, dizziness and h/o progressive deafness, which of the following is not a D/D:

Options :

- ✘ Acoustic neuroma
- ✘ Endolymphatic
- ✘ Meningitis
- ✔ Histiocytosis X

Question Number : 17 Question Id : 5799659669 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All of the following are true regarding Warthin's tumor except:

Options :

- ✔ More common in females
- ✘ Commonly involve the parotid glands
- ✘ They arise from the epithelial and the lymphoid cells
- ✘ 10% are bilateral

Question Number : 18 Question Id : 5799659670 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

CSF rhinorrhea is diagnosed by:

Options :

- ✘ Beta-2 microglobulin
- ✔ Beta-2 transferrin
- ✘ Thyroglobulin
- ✘ Transthyretin

Question Number : 19 Question Id : 5799659671 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The phrenic nerve:

Options :

- ✔ Runs from lateral to medial, over the scalenus anterior muscle, deep to the transverse cervical vessels.
- ✘ Runs from medial to lateral, over the scalenus anterior muscle, deep to the transverse cervical vessels.
- ✘ Runs from medial to lateral, over the scalenus anterior muscle, superficial to the transverse cervical vessels.
- ✘ Runs from lateral to medial, over the scalenus anterior muscle, and has a variable relationship with the transverse cervical vessels

Question Number : 20 Question Id : 5799659672 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The neck staging of a patient with nasopharyngeal SCC with a 4-cm metastatic node in the supraclavicular fossa on the right side and a 2-cm metastatic node on the left side at level II is:

Options :

1. ✘ N1a
2. ✘ N1b
3. ✘ N2c
4. ✔ N3

Question Number : 21 Question Id : 5799659673 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a boundary of "Chaussaignac triangle" in Neck dissection:

Options :

1. ✘ Subclavian artery as base
2. ✘ Laterally Longus colli and scalenus anterior attached to tubercle of C-6
3. ✘ Midline medially
4. ✔ Internal carotid artery laterally

Question Number : 22 Question Id : 5799659674 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In vestibular physiology, the Ewald's second law states that:

Options :

1. ✔ Excitatory responses are of higher amplitude than inhibitory response
2. ✘ Inhibitory responses are of higher amplitude than excitatory response
3. ✘ Mechanical stimulation of the membranous canal will lead to eye and head movements in the plane of that canal.
4. ✘ Mechanical stimulation of the membranous canal will lead to eye and head movements perpendicular to the plane of that canal.

Question Number : 23 Question Id : 5799659675 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following structure divides the Glasscock and Kawase triangle:

Options :

1. ✘ Internal Carotid Artery
2. ✔ Greater Superficial petrosal nerve
3. ✘ Deep Petrosal Nerve
4. ✘ Nerve to Pterygoid canal

Question Number : 24 Question Id : 5799659676 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False statement about acoustic rhinomanometry:

Options :

1. ✔ Inferior method as compared to active rhinomanometry
2. ✘ The plot of cross-sectional area against distance can also be expressed as nasal volume for given distances along the nasal passage.
3. ✘ it measures the nasal cross sectional area along with the length of nasal airway
4. ✘ sound pulse is used to measure nasal resistance

Question Number : 25 Question Id : 5799659677 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Interlamellar cell of Grunwald is defined as:

Options :

- the superior meatus may pneumatize the vertical lamella of the middle turbinate
- Inter frontal Cell
- pneumatized uncinata process
- pneumatized posterior ethmoidal cells extending to sphenoid sinus

Question Number : 26 Question Id : 5799659678 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is true regarding calyx endings in the vestibula neuroepithelium?

Options :

- Innervate type I and type II hair cells
- Are equally distributed throughout the epithelium
- Are seen only on calyx afferent neuron
- Receive postsynaptic afferent input

Question Number : 27 Question Id : 5799659679 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following anatomical features predisposes a rhinoplasty patient to postoperative nasal airway obstruction?

Options :

- Long nasal bones
- Wide upper cartilaginous vault
- Convex contour of the lower lateral cartilages
- Cephalically positioned lateral crura

Question Number : 28 Question Id : 5799659680 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A child with severe sensorineural hearing loss (SNHL) presents with fever, headache, nausea and vomiting, and photophobia a day after being diagnosed with acute otitis media (AOM). Temporal bone imaging is most likely to demonstrate:

Options :

- Dehiscence of the tegmen mastoideum
- Patent tympanomeningeal fissure
- Mondini malformation
- Congenital cholesteatoma

Question Number : 29 Question Id : 5799659681 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a contraindication to percutaneous tracheostomy?

Options :

- High innominate artery
- Body mass index (BMI) > 30

- ✘ Large midline neck mass
- ✘ Unprotected airway

Question Number : 30 Question Id : 5799659682 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is a true statement about complications following tonsillectomy?

Options :

- ✘ Secondary bleeding is more common than primary bleeding.
- ✘ In a child with submucous cleft palate, tonsillectomy is contraindicated.
- ✘ Children under 3 years are more likely to have respiratory complications.
- ✔ Secondary bleeding is more common than primary bleeding and Children under 3 years are more likely to have respiratory complications.

Question Number : 31 Question Id : 5799659683 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True about cilia of nasal mucosa are all except

Options :

- ✘ Have a 9+2 microtubular structure
- ✔ Beat at a frequency of approximately 120Hz
- ✘ May be functionally assessed by nasal brush biopsy
- ✘ May be functionally assessed by the saccharin test

Question Number : 32 Question Id : 5799659684 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

According to 2009 the Bethesda System for Reporting Thyroid Cytopathology, what is the risk of malignancy of follicular lesion of undetermined significance (FLUS)?

Options :

- ✔ 5% to 15%
- ✘ <5%
- ✘ 20% to 30%
- ✘ >50%

Question Number : 33 Question Id : 5799659685 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

What is not true about Prussack's space?

Options :

- ✘ pars flaccida cholesteatoma starts in this space
- ✘ lateral boundary is shrapnell's membrane
- ✔ Ventilated through tympanic isthmus
- ✘ the space can be seen radiologically on coronal oblique image

Question Number : 34 Question Id : 5799659686 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All are true about brachiocephalic vein except:

Options :

- ✘ two join to form SVC

- right brachiocephalic vein is longer than left brachiocephalic vein
- inferior thyroid vein is one of the contributors
- originate deep to left sternoclavicular joint

Question Number : 35 Question Id : 5799659687 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False statement in the following is:

Options :

- inferior thyroid veins maybe 1,2,3,4 in number
- colour Doppler of ITA is useful in assessing the functional state
- ITV may join the other side before ending into brachiocephalic vein
- None of the given options are correct

Question Number : 36 Question Id : 5799659688 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False about jugulodigastric lymph nodes is:

Options :

- significant only when size >1.5 cms radiologically
- can be palpable in normal individuals
- may get involved in infections of areas outside otorhinolaryngology
- chronic tonsillitis is the most common cause .

Question Number : 37 Question Id : 5799659689 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Brachio-oto-renal syndrome is NOT characterised by:

Options :

- autosomal dominant disorder
- can have middle and lower ear abnormality
- can have brachial cleft anomaly
- problem is with chromosome 9q arm

Question Number : 38 Question Id : 5799659690 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

What is wrong about suprasternal space?

Options :

- lies between two heads of SCM muscle with their enveloping fascia
- extend from sternum to half way up the neck
- contains fat & communicating vein between two anterior jugular veins
- contains LN

Question Number : 39 Question Id : 5799659691 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Sublingual glands drain via

Options :

- Stenson duct

- ✘ Wharton duct
- ✘ Bartholin duct
- ✔ Duct of Rivinus

Question Number : 40 Question Id : 579965962 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

HPV positive HNSCC has all except:

Options :

- ✘ occurs in non-smokers
- ✘ patients are relatively younger
- ✘ more radiosensitive
- ✔ p53 mutational loss common

Question Number : 41 Question Id : 579965963 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

TRUE statement about vestibular neuronitis is:

Options :

- ✘ viral in origin
- ✘ risk of developing BPPV later on
- ✘ steroids systemic or intratympanic are given
- ✔ All the given options are correct

Question Number : 42 Question Id : 579965964 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Desirable minimum close microscopic margin in oral tongue cancer is

Options :

- ✘ 1 to 5 mm
- ✘ 1 to 2 mm
- ✘ atleast 3 mm
- ✔ it depends upon the situation and very difficult to decide precise margin

Question Number : 43 Question Id : 579965965 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Expansile lytic osseous metastasis are characteristic of primary malignancy of:

Options :

- ✔ thyroid
- ✘ bronchus
- ✘ pinna
- ✘ tongue

Question Number : 44 Question Id : 579965966 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Sensorineural deafness is seen in A/E-

Options :

- ✘ Alport's syndrome

- ✘ Pendred's syndrome
- ✘ Treacher Collins syndrome
- ✔ Crouzon's syndrome

Question Number : 45 Question Id : 5799659697 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a histological variant of thyroid neoplasm-

Options :

- ✘ Follicular cell
- ✔ Merkel cell
- ✘ Insular cell
- ✘ Anaplastic

Question Number : 46 Question Id : 5799659698 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Radioiodine destroys the neoplastic cells predominantly by-

Options :

- ✘ X - rays
- ✔ beta rays
- ✘ gamma rays
- ✘ alpha particles

Question Number : 47 Question Id : 5799659699 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Contraindications of supraglottic laryngectomy are A/E:

Options :

- ✘ Poor pulmonary reserve
- ✘ Involvement of pyriform sinus
- ✔ Involvement of preepiglottic space
- ✘ Vocal cord fixation

Question Number : 48 Question Id : 5799659700 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A patient of head injury was brought to hospital, conscious, having clear nasal discharge through right nostril. NCCT head revealed non-operable injury to frontobasal area. Most appropriate management is:

Options :

- ✔ Wait & watch for 4- 5 days to allow spontaneous healing
- ✘ Do an MRI to localize the leak & control it endoscopically
- ✘ Put a dural catheter
- ✘ Approach transcranially to repair the damaged frontobasal region

Question Number : 49 Question Id : 5799659701 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True about Reinke's oedema:

Options :

1. ✘ Usually unilateral
2. ✔ Common In smoker
3. ✘ Corticosteroid is contraindicated
4. ✘ Involves a part of membranous vocal cords

Question Number : 50 Question Id : 5799659702 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Feature of Scheibe's syndrome is:

Options :

1. ✘ Semicircular canal fistula
2. ✘ Abnormality of bony labyrinth
3. ✔ Dysplasia of cochlea
4. ✘ Middle ear anomaly

Question Number : 51 Question Id : 5799659703 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True about nasopharyngeal carcinoma:

Options :

1. ✘ radiotherapy is treatment of choice
2. ✘ may be associated with unilateral otitis media
3. ✘ level 4 lymph nodes are not involved
4. ✔ All the given options are correct

Question Number : 52 Question Id : 5799659704 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Evidence based therapy of Bell's palsy include:

Options :

1. ✘ facial nerve massage
2. ✘ facial nerve stimulation
3. ✔ steroid
4. ✘ acyclovir

Question Number : 53 Question Id : 5799659705 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Person with vertigo without CNS involvement. Causes are:

Options :

1. ✘ perilymph fistula
2. ✘ otolithiasis
3. ✘ vestibular neuritis
4. ✔ All the given options are correct

Question Number : 54 Question Id : 5799659706 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True about tubercular otitis media:

Options :

1. ✓ pale granulation tissue in middle ear
2. ✗ no extra auricular complication
3. ✗ corticosteroids are used in treatment
4. ✗ painful

Question Number : 55 Question Id : 5799659707 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True about tympanometry:

Options :

1. ✗ flat in ossicular discontinuity
2. ✓ C type in Eustachian tube blockage
3. ✗ Ad type in otosclerosis
4. ✗ None of the given options are correct

Question Number : 56 Question Id : 5799659708 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True about bilateral abductor cord palsy

Options :

1. ✗ Voice is poor
2. ✓ Stridor is present
3. ✗ Urgent tracheostomy is needed
4. ✗ All the given options are correct

Question Number : 57 Question Id : 5799659709 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True about ranula is:

Options :

1. ✓ simple variety is true retention cyst
2. ✗ parotid is most common site
3. ✗ involves floor of mouth on both sides of frenulum
4. ✗ never extends into neck.

Question Number : 58 Question Id : 5799659710 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All are true about mandible fracture except:

Options :

1. ✗ malocclusion of teeth occur
2. ✓ anterior superior alveolar nerve is most commonly injured
3. ✗ condylar neck is most common site.
4. ✗ panorex radiographs are useful

Question Number : 59 Question Id : 5799659711 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Not a pernalignant lesion of oral cavity:

Options :

- ✘ leukoplakia
- ✘ lichen planus
- ✔ aphthous ulcer
- ✘ submucous fibrosis

Question Number : 60 Question Id : 5799659712 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True about grommet insertion:

Options :

- ✘ maximum duration of tube placement is 5 months.
- ✔ it is placed anteriorly on tympanic membrane.
- ✘ surgery is always needed to remove it.
- ✘ healing is slower after extrusion than after removal

Question Number : 61 Question Id : 5799659713 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Structure preserved in radical neck dissection

Options :

- ✘ internal jugular vein
- ✘ accessory nerve
- ✔ brachial plexus
- ✘ sternocleidomastoid muscle

Question Number : 62 Question Id : 5799659714 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is a scoring system for severity of wound infection & is particularly useful for surveillance & research-

Options :

- ✘ APGAR Score
- ✘ Glasgow scoring system
- ✔ Southampton grading system
- ✘ ASA classification

Question Number : 63 Question Id : 5799659715 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True about vestibular schwannoma

Options :

- ✘ more common in females
- ✘ second most common tumor of CP angle
- ✘ uncapsulated
- ✔ bilateral tumors seen in neurofibromatosis

Question Number : 64 Question Id : 5799659716 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Tensor of vocal cords is:

Options :

1. ✓ cricothyroid
2. ✗ interarytenoid
3. ✗ arytenoid
4. ✗ posterior cricoarytenoid

Question Number : 65 Question Id : 5799659717 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True about allergic fungal rhinosinusitis:

Options :

1. ✗ invasion of sinus mucosa with fungus
2. ✗ antifungal treatment leads to improvement
3. ✓ surgical clearance is mainstay
4. ✗ CT scan shows mucosal thickening with hypodense areas

Question Number : 66 Question Id : 5799659718 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Foreign body in trachea and bronchus can cause:

Options :

1. ✗ atelectasis
2. ✗ bronchiectasis
3. ✓ Both atelectasis and bronchiectasis
4. ✗ None of the given options are correct

Question Number : 67 Question Id : 5799659719 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True about conductive hearing loss:

Options :

1. ✗ speech discrimination score is poor
2. ✗ high frequencies are affected more
3. ✓ loss is not more than 60 db
4. ✗ congenital cholesteatoma is not cause

Question Number : 68 Question Id : 5799659720 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True about croup:

Options :

1. ✗ caused by H. influenza virus
2. ✓ X-ray chest shows steeple sign of subglottic narrowing
3. ✗ common age of presentation is more than 3 years.
4. ✗ stridor is absent

Question Number : 69 Question Id : 5799659721 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

FALSE about Mobius syndrome

Options :

1. ✘ 6th cranial nerve spared
2. ✘ abduction defect always bilateral
3. ✔ esotropia commonly seen
4. ✘ unilateral facial nerve involvement

Question Number : 70 Question Id : 5799659722 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common type of trachea-esophageal fistula is-

Options :

1. ✔ Proximal end blind, distal end communicating with trachea
2. ✘ Distal end blind, proximal end communicating with trachea
3. ✘ Both ends blind
4. ✘ Both ends open

Question Number : 71 Question Id : 5799659723 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

FALSE about acute epiglottitis:

Options :

1. ✘ prodromal symptoms are present
2. ✘ caused by parainfluenza virus type 1 and 2
3. ✘ thumb sign in X-ray
4. ✔ cough is usually absent

Question Number : 72 Question Id : 5799659724 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

FALSE about ethmoid sinus:

Options :

1. ✘ Not present at birth
2. ✔ their number varies from 3 -18
3. ✘ first radiological evidence at 3 years
4. ✘ reaches adult size by 6 years

Question Number : 73 Question Id : 5799659725 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Development in all parts of ear (ext+middle+int) is completed by ___weeks of IUL:

Options :

1. ✘ 16th week
2. ✘ 20th week
3. ✘ 28th week
4. ✔ 30th week

Question Number : 74 Question Id : 5799659726 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Isthmus in external auditory canal is _____from tympanic membrane

Options :

- ✘ 5 mm
- ✘ 3 mm
- ✘ 4 mm
- ✔ 6 mm

Question Number : 75 Question Id : 5799659727 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In a true lateral radiograph of the skull and in coronal section through the temporal bone external and internal acoustic meatus are superimposed. Best view for mastoid is

Options :

- ✔ Schuller's view
- ✘ Law's lateral view
- ✘ Law's lateral oblique view
- ✘ Towne's view

Question Number : 76 Question Id : 5799659728 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Following things about nasal intubation are not true-

Options :

- ✘ required where there is trismus
- ✘ Done where manipulation of neck is considered unsafe
- ✘ may require sedation in uncooperative patient
- ✔ easier to perform than orotracheal intubation and therefore no complication

Question Number : 77 Question Id : 5799659729 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The most common malignant salivary gland neoplasm is:

Options :

- ✔ Mucoepidermoid carcinoma
- ✘ Adenoid cystic carcinoma
- ✘ Pleomorphic adenoma
- ✘ Acinic cell carcinoma

Question Number : 78 Question Id : 5799659730 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The most common cause for inspiratory stridor in a neonate is:

Options :

- ✘ b/l vocal cord paralysis
- ✔ laryngomalacia
- ✘ congenital laryngeal web
- ✘ congenital subglottic stenosis

Question Number : 79 Question Id : 5799659731 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a feature of MEN-2:

Options :

1. ✘ Medulary carcinoma thyroid
2. ✘ Parathyroid adenoma
3. ✔ Pituitary Adenoma
4. ✘ Pheochromocytoma

Question Number : 80 Question Id : 5799659732 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Moh's Micrographic excision of Basal cell carcinoma is used for all of the following except:

Options :

1. ✘ Recurrent tumour
2. ✔ Tumour less than 2 cm in diameter
3. ✘ Tumour with aggressive histology
4. ✘ Tumour with perineural invasion

Question Number : 81 Question Id : 5799659733 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Dentigerous cyst arises from:

Options :

1. ✘ The root of a caries tooth
2. ✘ the periosteum of fractured mandible
3. ✔ An unerupted permanent tooth
4. ✘ The sequestrum of osteomyelitis of mandible

Question Number : 82 Question Id : 5799659734 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The first total laryngectomy was performed by:

Options :

1. ✘ Crile
2. ✔ Billroth
3. ✘ Kocher
4. ✘ Gluck Sorenson

Question Number : 83 Question Id : 5799659735 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following represents correct boundary of sinus tympani:

Options :

1. ✘ superiorly posterior semicircular canal
2. ✘ Medially subiculum
3. ✘ inferiorly ponticulus
4. ✔ Laterally facial nerve

Question Number : 84 Question Id : 5799659736 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following two LASERs can be used in Stapedotomy:

Options :

- ✘ KTP and CO₂ Laser
- ✔ CO₂ and Nd:YAG Laser
- ✘ Nd:YAG and KTP Laser
- ✘ He-Neon and CO₂ Laser

Question Number : 85 Question Id : 5799659737 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False statement regarding resistant otomycosis:

Options :

- ✔ topical antifungals are treatment of choice
- ✘ immunotherapy is the treatment of choice.
- ✘ it is required to exclude other fungal infections like athlete's foot
- ✘ mainly cause by dermatophytes

Question Number : 86 Question Id : 5799659738 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False statement regarding Ramsay Hunt Syndrome:

Options :

- ✘ 8th nerve may be involved leading to hearing loss, tinnitus and vertigo.
- ✘ In some cases, mucosal rash can be present.
- ✔ Reactivation of Herpes Zoster virus at second genu
- ✘ Auricular pain is usually first symptom

Question Number : 87 Question Id : 5799659739 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following nerve is not involved in Vernet Syndrome:

Options :

- ✔ 7th
- ✘ 9th
- ✘ 10th
- ✘ 11th

Question Number : 88 Question Id : 5799659740 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False statement regarding treatment options of recurrent generalised furunculosis:

Options :

- ✘ eradication therapy with nasal mupirocin
- ✘ eradication therapy with oral flucloxacillin for 14 days
- ✘ bacterial interference therapy
- ✔ eradication therapy with steroids

Question Number : 89 Question Id : 5799659741 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False statement regarding granular myringitis:

Options :

1. ✓ most common organism is Staphylococcal aureus
2. ✗ granulations present over tympanic membrane
3. ✗ most common symptom is foul smelling discharge
4. ✗ local debridement and topical antibiotic are treatment of choice

Question Number : 90 Question Id : 5799659742 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False statement about lupus vulgaris:

Options :

1. ✗ Causative organism are Mycobacterium tuberculosis and M.bovis
2. ✗ Apple Jelly nodules are chacterstic
3. ✓ Second line antitubercular drugs are required in all the cases
4. ✗ In a chronic lesion, malignancy can occur mainly Squamous, rarely BCC

Question Number : 91 Question Id : 5799659743 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a part of Pierre Robin Sequence:

Options :

1. ✓ Laryngomalacia
2. ✗ Retrognathia/Micrognathia
3. ✗ Glossoptosis
4. ✗ Cleft palate

Question Number : 92 Question Id : 5799659744 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False statement about Bath Plug Technique of CSF leak repair:

Options :

1. ✗ Most commonly used technique
2. ✗ Thedural defect is enlarged until the bony rim of the skullbase can be clearly seen
3. ✗ Fat graft is harvested from ear lobe
4. ✓ Fibrin glue should not be used for sealing

Question Number : 93 Question Id : 5799659745 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is suggestive of Kero's type2:

Options :

1. ✗ olfactory fossa depth 1-3mm
2. ✓ olfactory fossa depth 4-7mm
3. ✗ olfactory fossa depth 8-17mm
4. ✗ olfactory fossa depth >17mm

Question Number : 94 Question Id : 5799659746 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The approximate incidence of dehiscence of bony canal over optic nerve and internal carotid artery in sphenoid sinus is:

Options :

1. ✘ 25% and 6% respectively
2. ✔ 6% and 25% respectively
3. ✘ 30% and 10% respectively
4. ✘ 10% and 30% respectively

Question Number : 95 Question Id : 5799659747 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Not a self-retaining hand retractor is

Options :

1. ✘ Jansen's mastoid retractor
2. ✘ Davis retractor
3. ✔ Mollison's mastoid retractor
4. ✘ Lempert's endaural retractor

Question Number : 96 Question Id : 5799659748 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

FALSE about rhinoscleroma

Options :

1. ✘ caused by gram positive bacteria
2. ✘ dapsonsone has been tried with some success
3. ✔ streptomycin and tetracycline are given together
4. ✘ causative organism cannot be cultured

Question Number : 97 Question Id : 5799659749 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Minor factors in CRS, tick the wrong statement

Options :

1. ✘ headache
2. ✘ fever
3. ✘ dental pain
4. ✔ hyposmia

Question Number : 98 Question Id : 5799659750 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Following things about trigeminal ganglion incorrect-

Options :

1. ✔ it gives five branches -ophthalmic, maxillary, mandibular, ciliary and frontal
2. ✘ it is semilunar in shape
3. ✘ it lies on anterior surface of petrous part of temporal bone near the apex
4. ✘ close association with plexus on internal carotid artery

Question Number : 99 Question Id : 5799659751 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Bilateral involvement of lymph nodes/ tongue is there in cancer, which one is most important

Options :

- ✘ Tongue is a midline structure
- ✘ Tongue drains to submental lymphnodes which is in midline
- ✘ There is rich lymphatic capillary plexus in mucous membrane of tongue
- ✔ Superficial lymphatic plexus on right and left side of tongue cross and meet on opposite side

Question Number : 100 Question Id : 5799659752 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

false statement about mastoid antrum:

Options :

- ✘ always present at birth -persistent /consistent
- ✘ lies 15mm medial to suprameatal triangle
- ✘ much more superficial in a child (as superficial as 2mm)
- ✔ may be absent many times

Question Number : 101 Question Id : 5799659753 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

false statement about Otitis media externa:

Options :

- ✘ children present with hearing loss
- ✔ they never present with balance disorder
- ✘ role of usg (A-MODE)in detecting middle ear effusion
- ✘ selective IgG2,IgG4 deficiencies may predispose to repeated otitis media

Question Number : 102 Question Id : 5799659754 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

ENOG in facial nerve injury /paralysis will be representative of:

Options :

- ✘ Done within 24 hrs
- ✔ Done within 72 hrs
- ✘ Done within 1 week
- ✘ As soon as injury is noticed

Question Number : 103 Question Id : 5799659755 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

What is not true about malignant tumor of minor salivary gland:

Options :

- ✘ commoner than major salivary gland
- ✘ most common are adenoid cystic carcinoma
- ✘ most common site of origin is hard /soft palate
- ✔ perineural invasion is less common therefore surgery is successful

Question Number : 104 Question Id : 5799659756 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a feature of granulomatosis with polyangitis

Options :

1. ✘ nasal polyp
2. ✔ perforated nasal septum
3. ✘ lower respiratory tract is spared
4. ✘ non blood stained nasal crusts

Question Number : 105 Question Id : 5799659757 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The mechanism of nasal allergy is:

Options :

1. ✔ Type 1 hypersensitivity reaction.
2. ✘ Type 2 hypersensitivity reaction.
3. ✘ Type 3 hypersensitivity reaction.
4. ✘ Type 4 hypersensitivity reaction.

Question Number : 106 Question Id : 5799659758 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Blood loss during major surgery is best assessed by-

Options :

1. ✘ visual assessment
2. ✔ gravimetric method
3. ✘ USG Doppler
4. ✘ Cardiac output by thermodilution.

Question Number : 107 Question Id : 5799659759 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Direction of nasolacrimal duct is

Options :

1. ✘ downward, backward & medially
2. ✔ downward, backward & laterally
3. ✘ downward, forward & medially
4. ✘ downward, forward & laterally

Question Number : 108 Question Id : 5799659760 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common complication of CALDWELL LUC OPERATION is

Options :

1. ✘ OA fistula
2. ✔ Infraorbital nerve palsy
3. ✘ Haemorrhage
4. ✘ Orbital cellulitis

Question Number : 109 Question Id : 5799659761 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All are true about nasolabial cysts except-

Options :

1. ✘ They are bilateral
2. ✘ Present in adulthood
3. ✔ derived from odontogenic epithelium
4. ✘ strong female predilection

Question Number : 110 Question Id : 5799659762 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All are true about tracheostomy & larynx in children except-

Options :

1. ✘ Omega- shaped epiglottis
2. ✘ Larynx is high
3. ✔ Trachea can be easily palpated
4. ✘ avoid too much neck extension during positioning

Question Number : 111 Question Id : 5799659763 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The subdermal plexus forms the vascular basis for which flap-

Options :

1. ✔ Randomised
2. ✘ Axial
3. ✘ Mucocutaneous
4. ✘ Fasciocutaneous

Question Number : 112 Question Id : 5799659764 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A 36 yr old obese man suffering from hypertension & snoring. Patient is a known smoker. In sleep test, there were 5 apnea/hyperapnea episodes per hour. He was give antihypertensives& advised to quit smoking. Next line of management is-

Options :

1. ✘ Uvulopalatopharyngoplasty
2. ✔ Diet plan & weight reduction
3. ✘ Nasal CPAP
4. ✘ Mandibular repositioning sling

Question Number : 113 Question Id : 5799659765 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Best time for Antibiotic administration for surgery is-

Options :

1. ✘ 1 hour before surgery
2. ✘ i/m 6 hrs before surgery
3. ✔ At the time of induction of anaesthesia
4. ✘ Orally given the night before.

Question Number : 114 Question Id : 5799659766 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

cavernous sinus thrombosis, which statement is incorrect:

Options :

- ✘ conjunctival and eyelid oedema
- ✘ ophthalmoplegias
- ✘ meningitis and brain abscess signs and symptoms
- ✔ None of the given options are correct

Question Number : 115 Question Id : 5799659767 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

which of the following is not true about uncinat process:

Options :

- ✔ coronally oriented bone
- ✘ can get attached above lamina papyracea
- ✘ it is pneumatised sometimes
- ✘ it runs in a sickle shaped curve from anteroinferior to posterosuperior position

Question Number : 116 Question Id : 5799659768 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

which of the following is not a branch of vagus:

Options :

- ✘ meningeal branch
- ✘ auricular branch
- ✘ superior laryngeal nerve
- ✔ external laryngeal nerve

Question Number : 117 Question Id : 5799659769 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

FALSE statement regarding thyroid scan is:

Options :

- ✘ functional activity of thyroid
- ✘ done initially with RAI
- ✘ now with Tc 99 injected IV
- ✔ iodine scan not used now a days

Question Number : 118 Question Id : 5799659770 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

false about monoclonal antibody:

Options :

- ✘ all names end with Mab necessarily
- ✘ these can be made in laboratory
- ✘ can be given S/C or IV or oral
- ✔ no sideeffects as they block molecules that stop the immune system working

Question Number : 119 Question Id : 5799659771 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False statement about juvenile recurrent respiratory papillomatosis:

Options :

1. ✘ Caused by HPV6 and 11
2. ✘ Multiple papilloma seen in respiratory tract
3. ✘ Excision with powered microdebrider is the gold standard for laryngeal papilloma
4. ✔ CO₂ LASER should not be used as it leads to recurrence and thermal trauma

Question Number : 120 Question Id : 5799659772 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a part of DiGeorge anomaly:

Options :

1. ✘ Partial or complete aplasia of thyroid and parathyroid glands
2. ✘ Cardiac anomaly
3. ✔ Vertebral anomaly
4. ✘ Craniofacial anomaly

Question Number : 121 Question Id : 5799659773 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Type 3 frontal cell is defined as:

Options :

1. ✘ Single cell above the frontal recess
2. ✘ Loner cell
3. ✔ Cell extending into frontal recess but occupying less than 50% of frontal sinus
4. ✘ Cell extending into frontal recess but occupying more than 50% of frontal sinus

Question Number : 122 Question Id : 5799659774 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Dysphagia in cardiac achalasia is:

Options :

1. ✘ Progressive
2. ✘ Regressive
3. ✘ Stationary
4. ✔ Intermittent

Question Number : 123 Question Id : 5799659775 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Neurofibromatosis type 2 is associated with all except:

Options :

1. ✘ B/L Acoustic Neuroma
2. ✘ Café-au lait spots
3. ✘ Chromosome 22
4. ✔ Lisch nodule

Question Number : 124 Question Id : 5799659776 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Hitzelberger's sign is?

Options :

1. ✘ Reduced corneal sensitivity
2. ✔ Hypoesthesia of the posterior meatal wall
3. ✘ Paraesthesia of face
4. ✘ Delayed blink reflex

Question Number : 125 Question Id : 5799659777 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Orbital apex syndrome involves?

Options :

1. ✘ Olfactory nerve
2. ✘ Ophthalmic division of trigeminal nerve
3. ✔ Maxillary division of trigeminal nerve
4. ✘ Mandibular division of trigeminal nerve

Question Number : 126 Question Id : 5799659778 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Pain in tonsillar fossa and upper neck is characteristic of?

Options :

1. ✔ Eagle's syndrome
2. ✘ Apert's syndrome
3. ✘ Sickler's syndrome
4. ✘ Usher's syndrome

Question Number : 127 Question Id : 5799659779 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Ipsilateral immobility of soft palate is seen in?

Options :

1. ✘ Lermoyez syndrome
2. ✘ Ortner's syndrome
3. ✘ Costen's syndrome
4. ✔ Trotter's syndrome

Question Number : 128 Question Id : 5799659780 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The following is true for Gradenigo's syndrome?

Options :

1. ✘ Mastoid tenderness
2. ✔ Retro-orbital pain
3. ✘ Hearing loss
4. ✘ Sagging of posterosuperior meatal wall

Question Number : 129 Question Id : 5799659781 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which is the investigation of choice in assessing hearing loss in neonates?

Options :

1. ✘ Impedance audiometry
2. ✔ Brainstem Evoked Response Audiometry (BERA)
3. ✘ Free field audiometry
4. ✘ Behavioral audiometry

Question Number : 130 Question Id : 5799659782 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A 30 year old male is having Attic cholesteatoma of left ear with lateral sinus thrombophlebitis. Which of the following will be the operation of choice?

Options :

1. ✘ Intact canal wall mastoidectomy
2. ✘ Simple mastoidectomy with Tympanoplasty
3. ✔ Canal wall down mastoidectomy
4. ✘ Mastoidectomy with cavity obliteration

Question Number : 131 Question Id : 5799659783 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a typical feature of malignant otitis externa?

Options :

1. ✘ Caused by *Pseudomonas aeruginosa*
2. ✘ Patients are usually old
3. ✔ Mitotic figures are high
4. ✘ Patient is immune compromised

Question Number : 132 Question Id : 5799659784 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a typical feature of Ménière's disease?

Options :

1. ✘ Sensorineural deafness
2. ✔ Pulsatile tinnitus
3. ✘ Vertigo
4. ✘ Fluctuating deafness

Question Number : 133 Question Id : 5799659785 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A case of carcinoma larynx with the involvement of anterior commissure and right vocal cord, developed perichondritis of thyroid cartilage. Which of the following statements is true for the managements of this case?

Options :

1. ✘ He should be given radical radiotherapy as this can cure early tumours
2. ✘ He should be treated with combination of chemotherapy and radiotherapy
3. ✘ He should first receive radiotherapy and if residual tumour is present then should undergo laryngectomy
4. ✔ He should first undergo laryngectomy and then post—operative radiotherapy.

Question Number : 134 Question Id : 5799659786 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Hyperacusis in Bell's palsy is due to the the paralysis of the following muscle:

Options :

- ✘ Tensor tympani
- ✘ Levatorvelipalatini
- ✘ Tensor velipalatini
- ✔ Stapedius

Question Number : 135 Question Id : 5799659787 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is the most common etiological agent in paranasal sinus mycoses?

Options :

- ✔ Aspergillus spp
- ✘ Histoplasma
- ✘ Conidioboluscoronatus
- ✘ Candida albicans

Question Number : 136 Question Id : 5799659788 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Treatment of choice in a 6 year old child with recurrent URTI with mouth breathing and failure to grow with high arched palate and impaired hearing is :

Options :

- ✘ Tonsillectomy
- ✘ Grommet insertion
- ✘ Myringotomy with grommet insertion
- ✔ Adenoidectomy with grommet insertion

Question Number : 137 Question Id : 5799659789 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Treatment of choice for glue ear is

Options :

- ✘ Myringotomy with cold knife
- ✘ Myringotomy with diode laser
- ✔ Myringotomy with ventilation tube insertion
- ✘ Conservative treatment with analgesics & antibiotics

Question Number : 138 Question Id : 5799659790 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A lady has bilateral hearing loss since 4 years which worsened during pregnancy. Type of impedance audiometry graph will be :

Options :

- ✘ Ad
- ✔ As
- ✘ B

4. ✘ C

Question Number : 139 Question Id : 5799659791 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Rhinophyma is associated with:

Options :

1. ✔ Hypertrophy of the sebaceous glands
2. ✘ Hypertrophy of sweat glands
3. ✘ Hypertrophy of endothelial cells
4. ✘ Hypertrophy of epithelial cells

Question Number : 140 Question Id : 5799659792 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Rhinolalia clausa is associated with all of the following except?

Options :

1. ✘ Allergic rhinitis
2. ✔ Palatal paralysis
3. ✘ Adenoids
4. ✘ Nasal polyps

Question Number : 141 Question Id : 5799659793 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Gold standard test for laryngopharyngeal reflux is?

Options :

1. ✘ Esophageal motility study
2. ✘ Barium swallow
3. ✔ 24 hour double probe pH monitoring
4. ✘ Esophageal biopsy

Question Number : 142 Question Id : 5799659794 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Androphonia can be corrected by doing:

Options :

1. ✘ Type 1 Thyroplasty
2. ✘ Type 2 Thyroplasty
3. ✘ Type 3 Thyroplasty
4. ✔ Type 4 Thyroplasty

Question Number : 143 Question Id : 5799659795 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The most common and earliest manifestation of carcinoma of the glottis is:

Options :

1. ✔ Hoarseness
2. ✘ Haemoptysis
3. ✘ Cervical lymph nodes

4. ✘ Stridor

Question Number : 144 Question Id : 5799659796 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A 5 year old patient is scheduled for tonsillectomy. On the day of surgery he had running nose, temperature 37.5°C and dry cough. Which of the following should be the most appropriate decision for surgery?

Options :

1. ✘ Surgery should be cancelled
2. ✘ Can proceed for surgery if chest is clear and there is no history of asthma
3. ✘ Should get Xray chest before proceeding for surgery
4. ✔ Cancel surgery for 3 weeks and patient to be put on antibiotics

Question Number : 145 Question Id : 5799659797 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

During inspiration the main current of airflow in a normal nasal cavity is through:

Options :

1. ✔ Middle part of the cavity in middle meatus in a parabolic curve
2. ✘ Lower part of the cavity in the inferior meatus in a parabolic curve
3. ✘ Superior part of the cavity in the superior meatus
4. ✘ Through olfactory area

Question Number : 146 Question Id : 5799659798 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following conditions causes the maximum hearing loss?

Options :

1. ✔ Ossicular disruption with intact tympanic membrane
2. ✘ Disruption of malleus and incus as well tympanic membrane
3. ✘ Partial fixation of the stapes footplate
4. ✘ Otitis media with effusion

Question Number : 147 Question Id : 5799659799 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Ceruminous glands present in the Ear are ?

Options :

1. ✘ Modified Eccrine glands
2. ✔ Modified apocrine glands
3. ✘ Modified holocrine glands
4. ✘ Mucous glands

Question Number : 148 Question Id : 5799659800 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In lesion of facial nerve at vertical part "below the nerve of stapedius" there is:

Options :

1. ✔ Loss of taste
2. ✘ Impairment of salivation

3. ✘ Impairment of lacrimation
4. ✘ Hyperacusis

Question Number : 149 Question Id : 5799659801 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Valvular obstructive emphysema occurs due to:

Options :

1. ✔ Partial obstruction of the bronchus.
2. ✘ Complete obstruction of the bronchus.
3. ✘ When the pretracheal fascia is sutured tightly.
4. ✘ Injury of the pleura.

Question Number : 150 Question Id : 5799659802 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The early symptom of Bell's palsy is:

Options :

1. ✘ Dropping of angle of the affected side
2. ✘ Obliteration of the angle of the mouth
3. ✔ Pain of acute onset behind the ear
4. ✘ Inability to close the eye