

# Question Paper Preview

## Notations :

- Options shown in green color and with ✓ icon are correct.
- Options shown in red color and with ✗ icon are incorrect.

<b>Question Paper Name:</b>	Civil Assistant Surgeon in General Surgery 5th Dec 2017 Shift 1
<b>Subject Name:</b>	Civil Assistant Surgeon in General Surgery
<b>Creation Date:</b>	2017-12-05 14:04:27
<b>Duration:</b>	150
<b>Total Marks:</b>	150
<b>Display Marks:</b>	No
<b>Calculator:</b>	Scientific
<b>Magnifying Glass Required?:</b>	No
<b>Ruler Required?:</b>	No
<b>Eraser Required?:</b>	No
<b>Scratch Pad Required?:</b>	No
<b>Rough Sketch/Notepad Required?:</b>	No
<b>Protractor Required?:</b>	No

## Civil Assistant Surgeon in General Surgery

<b>Group Number :</b>	1
<b>Group Id :</b>	57996567
<b>Group Maximum Duration :</b>	0
<b>Group Minimum Duration :</b>	150
<b>Revisit allowed for view? :</b>	No
<b>Revisit allowed for edit? :</b>	No
<b>Break time:</b>	0
<b>Group Marks:</b>	150

## Civil Assistant Surgeon in General Surgery

<b>Section Id :</b>	57996567
<b>Section Number :</b>	1
<b>Section type :</b>	Online
<b>Mandatory or Optional:</b>	Mandatory
<b>Number of Questions:</b>	150
<b>Number of Questions to be attempted:</b>	150
<b>Section Marks:</b>	150
<b>Display Number Panel:</b>	Yes
<b>Group All Questions:</b>	No

<b>Sub-Section Number:</b>	1
<b>Sub-Section Id:</b>	579965105
<b>Question Shuffling Allowed :</b>	Yes

Question Number : 1 Question Id : 5799659953 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False regarding caecal volvulus:

Options :

1. ✘ Present with small bowel obstruction
2. ✘ Present with air fluid levels in right upper quadrant and convexity towards left
3. ✘ Endoscopic derotation is not effective like sigmoid volvulus
4. ✔ Caecopexy can be a form of treatment

Question Number : 2 Question Id : 5799659954 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not true about haemorrhoids?

Options :

1. ✘ Pruritus is not common
2. ✔ Can be palpated on digital rectal examination in absence of complications
3. ✘ Band ligation is most commonly done office procedure
4. ✘ None of the given options are correct

Question Number : 3 Question Id : 5799659955 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A 10 mm calculus in the right lower ureter associated with proximal hydronephrosis is best treated with:

Options :

1. ✘ ESWL
2. ✘ Antegrade percutaneous access
3. ✘ Open ureterolithotomy
4. ✔ Ureteroscopic retrieval

Question Number : 4 Question Id : 5799659956 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Renal tuberculosis originates in the:

Options :

1. ✔ Renal papilla
2. ✘ Renal medulla
3. ✘ Afferent tubules
4. ✘ Efferent arteriole of glomerulus

Question Number : 5 Question Id : 5799659957 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Acute graft rejection occurs within

Options :

1. ✘ Seconds
2. ✘ Minutes to days
3. ✔ Weeks to months
4. ✘ Months to years

Question Number : 6 Question Id : 5799659958 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

For kidney function the investigation of choice is

Options :

1. ✘ Ultrasound
2. ✘ IVP
3. ✔ DTPA Scan
4. ✘ CECT

Question Number : 7 Question Id : 5799659959 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Hypoxia is defined as oxygen saturation of less than:

Options :

1. ✘ 100 Percent
2. ✘ 95 Percent
3. ✔ 90 Percent
4. ✘ 85 Percent

Question Number : 8 Question Id : 5799659960 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Laparotomy in blunt trauma should be done in

Options :

1. ✔ Peritoneal air on imaging
2. ✘ Grade II spleen damage
3. ✘ Grade II liver damage
4. ✘ Mild haemoperitoneum

Question Number : 9 Question Id : 5799659961 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Treatment of choice in a patient who has developed massive left sided haemothorax within an hour of road traffic accident is:

Options :

1. ✘ Strapping of chest
2. ✘ Conservative, wait and watch
3. ✘ Endotracheal intubation + IPPV
4. ✔ Tube thoracostomy

Question Number : 10 Question Id : 5799659962 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Metastatic node of left axilla in a patient with Gastric tumor is called-

Options :

1. ✘ Virchow node
2. ✘ Sister mary joseph node
3. ✘ Rotter 's node

4. ✓ Irish node

Question Number : 11 Question Id : 5799659963 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is true about intussusception?

Options :

1. ✗ Common in neonates
2. ✗ Fever always present
3. ✗ Not associated with tumors of intestine
4. ✓ May be relieved by barium enema

Question Number : 12 Question Id : 5799659964 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common presentation of GERD

Options :

1. ✗ Belching
2. ✗ Vomiting
3. ✓ Heartburn
4. ✗ Bloating

Question Number : 13 Question Id : 5799659965 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Absolute contraindication to varicose vein surgery is

Options :

1. ✗ Varicose ulcer
2. ✓ Deep Vein thrombosis
3. ✗ Thrombophlebitis
4. ✗ Lipodermatosclerosis

Question Number : 14 Question Id : 5799659966 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A 50 years old patient presented with progressive jaundice. Liver function test was done in which conjugated serum bilirubin- 4. 8% and total bilirubin- 6. 7%, alkaline phosphatase- 550 IU, SGOT-50, SGPT-65. Most probable diagnosis is:

Options :

1. ✗ Jaundice due to choledocholithiasis
2. ✗ Dubin-Johnson syndrome
3. ✗ Viral hepatitis
4. ✓ Malignant obstructive jaundice

Question Number : 15 Question Id : 5799659967 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A chronic alcoholic presents with abdominal pain radiating to the back that responds to analgesics. At evaluation the pancreatic duct was found to be dilated and stones were noted in the tail of pancreas. The most appropriate management is:

Options :

1. ✘ Pancreatic tail resection
2. ✘ Pancreaticojejunostomy
3. ✘ Percutaneous removal of stone
4. ✔ Medical management

Question Number : 16 Question Id : 5799659968 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A 25-years old female presents with pyrexia for ten days, develops acute pain in periumblical region spreading all over the abdomen. What would be the most likely cause?

Options :

1. ✘ Perforation peritonitis due to intestinal tuberculosis
2. ✘ Generalized peritonitis due to appendicular perforation
3. ✔ Typhoid enteric perforation and peritonitis
4. ✘ Acute salpingo-oophoritis with peritonitis

Question Number : 17 Question Id : 5799659969 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A women of 35-years, comes to emergency department with symptoms of pain in abdomen and bilious vomiting but no distension of bowel. Abdominal X-ray showed no air fluid level. Diagnosis is:

Options :

1. ✘ Carcinoma rectum
2. ✔ Duodenal obstruction
3. ✘ Adynamic ileus
4. ✘ Pseudo-obstruction

Question Number : 18 Question Id : 5799659970 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A man aged 60 years has history of IHD and atherosclerosis. He presents with abdominal pain and maroon stools. Most likely diagnosis:

Options :

1. ✘ Acute intestinal obstruction
2. ✔ Acute mesenteric ischemia
3. ✘ Peritonitis
4. ✘ Appendicitis

Question Number : 19 Question Id : 5799659971 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Prolonged post-op ileus is best treated by:

Options :

1. ✘ Long tube insertion
2. ✘ Calcium pantothenate
3. ✔ Laparotomy and exploration
4. ✘ Peristaltic stimulants

Question Number : 20 Question Id : 5799659972 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A pro-inflammatory cytokine is

Options :

1. ✘ IL-4
2. ✘ IL-10
3. ✘ TGF- $\beta$
4. ✔ TNF

Question Number : 21 Question Id : 5799659973 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Criteria for diagnosis of ARDS includes

Options :

1. ✘ Cardiogenic pulmonary odema
2. ✘ Unilateral pulmonary infiltrates
3. ✘ PaO<sub>2</sub> /FiO<sub>2</sub> 201 to 300
4. ✔ PAWP < 18 mm of Hg

Question Number : 22 Question Id : 5799659974 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Intraabdominal pressure in grade II abdominal compartment syndrome

Options :

1. ✘ 21-25 mm of Hg
2. ✘ 10-14 mm of Hg
3. ✔ 15-20 mm of Hg
4. ✘ 25-35 mm of Hg

Question Number : 23 Question Id : 5799659975 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most probable diagnosis in case of a patient presenting with raised JVP, CVP 16 mm of Hg and persistent hypotension is

Options :

1. ✘ Pneumothorax
2. ✘ Splenic trauma
3. ✘ Head injury
4. ✔ Cardiac tamponade

Question Number : 24 Question Id : 5799659976 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

CSF rhinorrhea is found in

Options :

1. ✘ LE FORT type 1
2. ✔ LE FORT type 3
3. ✘ fracture zygomatic bone
4. ✘ fracture nasal bone

Question Number : 25 Question Id : 5799659977 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

SIRS includes all except

Options :

1. ✘ Hyperthermia (> 38\* C)
2. ✔ Tachycardia (> 90 /min ,on Beta Blockers)
3. ✘ Tachypnoea (>20 / min)
4. ✘ TLC (>12000/ul)

Question Number : 26 Question Id : 5799659978 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All are high risk premalignant lesions except-

Options :

1. ✘ Erythroplakia
2. ✘ Speckled erythroplakia
3. ✘ Chronic hyperplastic candidiasis
4. ✔ Oral sub mucus fibrosis

Question Number : 27 Question Id : 5799659979 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Conley's pointer is helpful in-

Options :

1. ✘ Thyroid surgery
2. ✘ Breast surgery
3. ✔ Parotid surgery
4. ✘ None of the given options are correct

Question Number : 28 Question Id : 5799659980 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In case of choledochal cyst , bile diversion in to the small intestine is not done because of risk of

Options :

1. ✘ Malignancy
2. ✘ Pancreatitis
3. ✔ Recurrent cholangitis
4. ✘ Gall stone

Question Number : 29 Question Id : 5799659981 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Bariatric operation without restrictive mechanism of action

Options :

1. ✔ Duodenal switch
2. ✘ Vertical banded gastroplasty
3. ✘ Laparoscopic sleeve gastrectomy
4. ✘ Roux-en-Y gastric bypass

Question Number : 30 Question Id : 5799659982 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False statement about Warthin's tumor

Options :

1. ✘ common in old age
2. ✘ appear as hot spot on radionuclide scan
3. ✘ superficial parotidectomy is sufficient
4. ✔ high malignant potential

Question Number : 31 Question Id : 5799659983 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All are true regarding Caroli's disease except?

Options :

1. ✘ Intrahepatic stone formation
2. ✘ Biliary stasis
3. ✔ Commonly presents with ascites
4. ✘ Malignant change can occur

Question Number : 32 Question Id : 5799659984 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All of the following are included in Mangled Extremity Severity Score (MESS) except the following?

Options :

1. ✘ Neurogenic injury
2. ✔ Energy of injury
3. ✘ Shock
4. ✘ Ischemia

Question Number : 33 Question Id : 5799659985 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Mandible fracture is found most commonly in relation to which tooth

Options :

1. ✘ central incisor
2. ✔ canines
3. ✘ premolar
4. ✘ molars

Question Number : 34 Question Id : 5799659986 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Aneurysm of popliteal artery should be treated if size is above

Options :

1. ✘ 1.0 cm
2. ✘ 1.5 cm
3. ✘ 2.0 cm
4. ✔ 2.5 cm

Question Number : 35 Question Id : 5799659987 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical



Correct : 1 Wrong : 0.33

Heerfordt's syndrome is sarcoidosis involving all except -

Options :

1. ✘ Parotid swelling
2. ✘ Anterior uveitis
3. ✘ Facial palsy
4. ✔ Posterior uveitis

Question Number : 36 Question Id : 5799659988 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common symptom of chronic pancreatitis is:

Options :

1. ✔ Abdominal pain
2. ✘ Cachexia
3. ✘ Weight loss
4. ✘ Steatorrhea

Question Number : 37 Question Id : 5799659989 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A 20-years old football player received a hard kick in the epigastrium. A large cystic swelling appeared in the epigastrium two weeks later. The most likely diagnosis is:

Options :

1. ✘ Hydatid cyst of liver
2. ✘ Amoebic liver abscess
3. ✔ Pseudopancreatic cyst
4. ✘ Hematoma of rectus sheath

Question Number : 38 Question Id : 5799659990 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Ramu is 60 years old male with Carcinoma descending colon presents with acute intestinal obstruction. In emergency department treatment of choice is:

Options :

1. ✘ Defunctioning colostomy
2. ✔ Hartmann's procedure
3. ✘ Total colectomy
4. ✘ Left hemicolectomy

Question Number : 39 Question Id : 5799659991 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A 25-years old patient presented with mass in right iliac fossa, which after laparotomy was found to be carcinoid of 2.5 cm in diameter. What will be next step in management?

Options :

1. ✘ Segmental resection
2. ✘ Appendectomy
3. ✔ Right hemicolectomy
4. ✘ Do yearly 5-HIAA assay

Question Number : 40 Question Id : 5799659992 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Prognosis for carcinoma rectum is best assessed by:

Options :

- ✘ Site of tumour
- ✔ Histological grading
- ✘ Size of tumors
- ✘ Duration of the symptoms

Question Number : 41 Question Id : 5799659993 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

What is true about carcinoma penis?

Options :

- ✘ Metastasis is rare
- ✘ Occurs more commonly in circumcised male
- ✔ Arises from corona of glans
- ✘ Pain is frequent

Question Number : 42 Question Id : 5799659994 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

With reference to varicocele, which one of the following is not true of it?

Options :

- ✔ Varicosity of cremastic veins
- ✘ Left side is affected usually
- ✘ Feel like a bag of worms
- ✘ May lead to infertility

Question Number : 43 Question Id : 5799659995 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In differential diagnosis of epididymo-orchitis and torsion it is important that:

Options :

- ✘ Elevation of testis in torsion relieves pain
- ✔ Elevation of testis in epididymo-orchitis relieves pain
- ✘ Tenderness is characteristic of torsion of testis
- ✘ None of the given options are correct

Question Number : 44 Question Id : 5799659996 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In a blast injury, which of the following organ is least vulnerable to the blast wave?

Options :

- ✘ G.I tract
- ✘ Lungs
- ✔ Liver
- ✘ Ear drum

Question Number : 45 Question Id : 5799659997 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Nonabsorbable suture is

Options :

1. ✘ Poliglecaprone
2. ✘ Polydioxanone
3. ✔ Polyester
4. ✘ Polyglactin

Question Number : 46 Question Id : 5799659998 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Formula for fluid resuscitation in pediatric patients is

Options :

1. ✘ Parkland formula
2. ✘ Brooke formula
3. ✔ Galveston formula
4. ✘ Muir and Barclay formula

Question Number : 47 Question Id : 5799659999 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common early cause of death in thermal burn injury is

Options :

1. ✔ Hypovolemia
2. ✘ Sepsis
3. ✘ Asphyxia
4. ✘ Associated trauma

Question Number : 48 Question Id : 57996510000 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Abdominal approach is used in rectal prolapse ?

Options :

1. ✘ Thiersch operation
2. ✘ Delorme's operation
3. ✘ Altemier's procedure
4. ✔ Ripstein's operation

Question Number : 49 Question Id : 57996510001 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a component of Total Parenteral Nutrition?

Options :

1. ✘ Amino acids
2. ✘ Fats
3. ✘ Vitamins
4. ✔ Fiber

Question Number : 50 Question Id : 57996510002 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which investigation is gold standard in investigation of upper GIT

Options :

1. ✘ CECT
2. ✘ MRI
3. ✘ Ultrasound
4. ✔ Flexible endoscopy

Question Number : 51 Question Id : 57996510003 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The line of management in a case of CSF rhinorrhoea :

Options :

1. ✘ Immediate surgery
2. ✘ Nasal packing only
3. ✔ conservative treatment with antibiotics and observation
4. ✘ X-ray skull and nasal packing

Question Number : 52 Question Id : 57996510004 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In the immediate post-operative period, changes in the level of potassium occurs due to

Options :

1. ✔ excessive excretion
2. ✘ Exchange with magnesium
3. ✘ Exchange with calcium
4. ✘ Retention in body

Question Number : 53 Question Id : 57996510005 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Shifting dullness is associated with

Options :

1. ✔ ascites
2. ✘ perforation of hollow viscera
3. ✘ interloop abscess
4. ✘ subphrenic abscess

Question Number : 54 Question Id : 57996510006 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Condition which builds within hemithorax resulting in col-lapsed lung, flattened diaphragm, contralateral mediastinal shift and compromised venous return to right side of heart is known as:

Options :

1. ✘ Open pneumothorax
2. ✘ Flail chest
3. ✘ Massive pulmonary hemorrhage

4. ✓ Tension pneumothorax

Question Number : 55 Question Id : 57996510007 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Quadrante lobe of liver is present between:

Options :

1. ✓ Groove for ligamentum teres and gallbladder
2. ✗ Inferior vena cava and fissure for ligamentum venosum
3. ✗ Groove for inferior vena cava and fissure for ligamentum venosum
4. ✗ Porta hepatis and falciform ligament

Question Number : 56 Question Id : 57996510008 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A Gallstone gets impacted most commonly in which part of common bile duct?

Options :

1. ✓ Supraduodenal
2. ✗ Retroduodenal
3. ✗ Ampulla of Vater
4. ✗ Common hepatic duct

Question Number : 57 Question Id : 57996510009 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Not a predisposing factor for carcinoma esophagus:

Options :

1. ✗ Diverticula
2. ✗ Human papilloma virus
3. ✓ Mediastinal fibrosis
4. ✗ Caustic ingestion

Question Number : 58 Question Id : 57996510010 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Early stage of carcinoma esophagus is diagnosed by:

Options :

1. ✗ Barium meal
2. ✓ Trans-esophageal USG
3. ✗ MRI
4. ✗ Endoscopy

Question Number : 59 Question Id : 57996510011 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The following statement about pilonidal sinus is true:

Options :

1. ✗ More common in females
2. ✗ Mostly congenital
3. ✗ Prognosis after surgery is poor

4. ✓ Treatment of choice is surgical excision of sinus tract

Question Number : 60 Question Id : 57996510012 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

DTPA scan of hypertensive young lady is normal, USG shows small kidney on left side. Next investigation will be:

Options :

1. ✗ CT Scan abdomen
2. ✗ Retrograde pyelogram
3. ✓ Digital subtraction angiography
4. ✗ None of the given options are correct

Question Number : 61 Question Id : 57996510013 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Not a feature of melanoma

Options :

1. ✗ UV Radiation is a key causative factor
2. ✗ Excisional biopsy is best method to diagnose
3. ✗ Superficial spreading type is most common
4. ✓ Radiosensitive tumor

Question Number : 62 Question Id : 57996510014 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Not a feature of phyllode tumor

Options :

1. ✗ Mixture of connective and epithelial tissues
2. ✗ Clusters of leaflike structures on histology
3. ✓ Diagnosed by cytology
4. ✗ Metastasis along hematogenous route

Question Number : 63 Question Id : 57996510015 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Portal vein is formed by

Options :

1. ✗ Inferior mesenteric vein & splenic vein
2. ✓ Sup mesenteric vein & splenic vein
3. ✗ Inferior & superior mesenteric vein
4. ✗ Splenic vein inf& sup mesenteric vein

Question Number : 64 Question Id : 57996510016 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Lord's plication is used in the treatment of ?

Options :

1. ✓ Hydrocele
2. ✗ Inguinal hernia

- ✘ Testicular malignancy
- ✘ Varicocele

Question Number : 65 Question Id : 57996510017 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following renal calculi is seen associated with proteus infection?

Options :

- ✘ Calcium Oxalate
- ✔ Triple Phosphate
- ✘ Xanthine
- ✘ Uric Acid

Question Number : 66 Question Id : 57996510018 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Lumbar sympathectomy is of value in the management of:

Options :

- ✘ Intermittent claudication
- ✔ Distal ischaemia affecting the skin of the toes
- ✘ Arteriovenous Fistula
- ✘ Back pain

Question Number : 67 Question Id : 57996510019 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is truly defunctional colostomy

Options :

- ✘ loop colostomy
- ✘ end colostomy
- ✔ double barrel colostomy
- ✘ None of the given options are correct

Question Number : 68 Question Id : 57996510020 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

One of the following is not included In case of airway clearance:

Options :

- ✘ Mouth gag
- ✔ Head lift
- ✘ Chin lift
- ✘ Neck tilt

Question Number : 69 Question Id : 57996510021 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A 34 year male presenting with apparently superficial injury to neck was brought dead in the emergency department due to road traffic accident sustained 10 minutes before presentation, probable cause of sudden death in this case is:

Options :

- ✔ Air embolism through superficial neck veins

- ✘ Injury to trachea
- ✘ Bleeding from subclavian artery
- ✘ Injury to phrenic nerve

Question Number : 70 Question Id : 57996510022 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Not a complication of laparoscopy:

Options :

- ✘ Diaphragmatic rupture
- ✘ Vascular injury
- ✘ Pneumothorax
- ✔ Pneumomediastinum

Question Number : 71 Question Id : 57996510023 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Centre for Diseases control (CDC) defines surgical site infection for non-prosthetic surgery up to :

Options :

- ✔ 30 days
- ✘ 90 days
- ✘ 180 days
- ✘ 1 year

Question Number : 72 Question Id : 57996510024 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Peritoneal dissemination of gastric cancer are best detected by-

Options :

- ✘ Abdominal ultrasound
- ✔ Laparoscopy
- ✘ Computed tomography
- ✘ MRI

Question Number : 73 Question Id : 57996510025 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Pyoderma Gangrenosum is most commonly associated with-

Options :

- ✔ Ulcerative colitis
- ✘ Crohn's disease
- ✘ Amoebic colitis
- ✘ Ischaemic colitis

Question Number : 74 Question Id : 57996510026 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common complication haemorrhoidectomy is

Options :

- ✘ Fecal incontinence



2. ✘ Infection
3. ✘ Haemorrhage
4. ✔ Urinary retention

Question Number : 75 Question Id : 57996510027 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Content of Amyand Hernia is

Options :

1. ✘ Caecum
2. ✔ Appendix
3. ✘ Urinary bladder
4. ✘ Sigmoid colon

Question Number : 76 Question Id : 57996510028 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A forty year old multipara female presented with true cut biopsy proved infiltrating ductal carcinoma in left breast measuring 4 cm in greatest diameter without skin and chest wall involvement. Patient also had USG and FNAB proved single mobile level II axillary lymph node involvement in right side. There was no other lymph node involvement or any other metastasis. What is the TNM staging of the disease according to AJCC ?

Options :

1. ✘ Stage IB
2. ✘ Stage IIB
3. ✘ Stage IIIB
4. ✔ Stage IV

Question Number : 77 Question Id : 57996510029 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Marker for Medullary carcinoma of thyroid is

Options :

1. ✔ Calcitonin
2. ✘ Thyroglobulin
3. ✘ Normetanephrine
4. ✘ Chromogranin A

Question Number : 78 Question Id : 57996510030 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

'Pie in the sky' prostate is found in

Options :

1. ✘ Benign hyperplasia of prostate
2. ✘ Carcinoma prostate
3. ✔ Rupture membranous urethra
4. ✘ Tubercular infection of prostate

Question Number : 79 Question Id : 57996510031 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

'golf hole ' ureter is seen in

Options :

1. ✘ Ureteric calculus
2. ✘ Ureteric polyp
3. ✔ Tuberculosis of ureter
4. ✘ Retroperitoneal fibrosis

Question Number : 80 Question Id : 57996510032 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Gold standard to diagnose pulmonary embolism

Options :

1. ✘ Chest X-ray
2. ✔ Pulmonary angiography
3. ✘ Ventilation perfusion scintiscan
4. ✘ CT scan chest

Question Number : 81 Question Id : 57996510033 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following vessel is commonly involved in extradural hematoma?

Options :

1. ✔ Middle meningeal artery
2. ✘ Venous sinuses
3. ✘ Bridging veins
4. ✘ Middle cerebral artery

Question Number : 82 Question Id : 57996510034 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common route of infection in kidney tuberculosis is

Options :

1. ✘ Ascending spread
2. ✘ Lymphatic spread
3. ✘ Direct invasion
4. ✔ Hematogenous

Question Number : 83 Question Id : 57996510035 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a selection criteria to day care surgery?

Options :

1. ✔ Age < 65 years
2. ✘ ASA (American Society of Anesthesiologists) class II
3. ✘ BMI < 40
4. ✘ Operation duration < 2 hours

Question Number : 84 Question Id : 57996510036 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

According to Le Fort classification of facial injury which class does not involve orbit?

**Options :**

1. ✓ Class I
2. ✗ Class II
3. ✗ Class III
4. ✗ All classes involve orbit

**Question Number : 85 Question Id : 57996510037 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical**

**Correct : 1 Wrong : 0.33**

Which of the following is not associated with Prune-Belly syndrome?

**Options :**

1. ✗ Lax lower abdominal wall
2. ✗ Dilated ureter
3. ✗ Undescended testis
4. ✓ Spina bifida

**Question Number : 86 Question Id : 57996510038 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical**

**Correct : 1 Wrong : 0.33**

Which of the following is not a method of femoral hernia repair?

**Options :**

1. ✓ Kelly's repair
2. ✗ Lockwood repair
3. ✗ Lotheissen repair
4. ✗ Mc Evedy repair

**Question Number : 87 Question Id : 57996510039 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical**

**Correct : 1 Wrong : 0.33**

Sportsman hernia is due to

**Options :**

1. ✗ Rupture of ligament of Cooper
2. ✓ Pubic symphysis diastasis
3. ✗ Widening of inguinal canal
4. ✗ Muscle hematoma

**Question Number : 88 Question Id : 57996510040 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical**

**Correct : 1 Wrong : 0.33**

Which of the following is not included in Cushing's triad?

**Options :**

1. ✗ Hypertension
2. ✗ Bradycardia
3. ✓ Hypotension
4. ✗ Irregular respiration

**Question Number : 89 Question Id : 57996510041 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical**

**Correct : 1 Wrong : 0.33**

Up to how many years after cholecystectomy, retained CBD stone can be found?

Options :

- ✘ 12 months
- ✘ 15 months
- ✘ 18 months
- ✔ 24 months

Question Number : 90 Question Id : 57996510042 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not associated with MEN 1 syndrome?

Options :

- ✔ Medullary carcinoma of thyroid
- ✘ Pancreatic tumour
- ✘ Parathyroid hyperplasia
- ✘ Pituitary adenoma

Question Number : 91 Question Id : 57996510043 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a capsular plate of liver?

Options :

- ✔ Portal plate
- ✘ Hilar plate
- ✘ Umbilical plate
- ✘ Cystic plate

Question Number : 92 Question Id : 57996510044 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Ray's amputation is

Options :

- ✔ Amputation of toe
- ✘ Amputation at the level of tarsal bones
- ✘ Amputation at the level of knee
- ✘ Amputation above knee

Question Number : 93 Question Id : 57996510045 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not removed in Auschincloss Mastectomy?

Options :

- ✘ Whole breast
- ✔ Pectoralis minor muscle
- ✘ Central group of axillary lymph node
- ✘ All of the given options are correct

Question Number : 94 Question Id : 57996510046 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not true about acute appendicitis?

Options :

1. ✘ Leukocytosis, often with a left shift of neutrophils is present in 90% cases.
2. ✔ C-reactive protein has high sensitivity but low specificity in diagnosis.
3. ✘ Pregnancy test is mandatory in women of childbearing age.
4. ✘ Intravenous contrast enhanced CT scan is recommended as an important diagnostic tool.

Question Number : 95 Question Id : 57996510047 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Chemotherapeutic agent of choice for anal canal carcinoma is

Options :

1. ✔ Mitomycin C
2. ✘ Cyclophosphamide
3. ✘ Vincristine
4. ✘ Bleomycin

Question Number : 96 Question Id : 57996510048 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a surgical treatment for rectal prolapsed?

Options :

1. ✘ Altemeier procedure
2. ✘ Goodsall's operation
3. ✘ Ripstein repair
4. ✔ Ferguson's procedure

Question Number : 97 Question Id : 57996510049 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Investigation of choice for posterior urethral valve is

Options :

1. ✘ RGU
2. ✔ MCU
3. ✘ IVU
4. ✘ USG

Question Number : 98 Question Id : 57996510050 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which is the most common site of primary malignancy for metastatic tumors of thyroid gland?

Options :

1. ✔ Breast
2. ✘ Lung
3. ✘ Tonsil
4. ✘ Liver

Question Number : 99 Question Id : 57996510051 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Deneyer shunt is used for

Options :

1. ✓ Ascitis
2. ✗ Dialysis
3. ✗ Raised ICT
4. ✗ Raised IOP

Question Number : 100 Question Id : 57996510052 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following sarcoma metastasis in lymph node?

Options :

1. ✓ Rhabdomyosarcoma
2. ✗ Osteosarcoma
3. ✗ Leiomyosarcoma
4. ✗ Fibrosarcoma

Question Number : 101 Question Id : 57996510053 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a component of MELD scoring for end stage liver disease?

Options :

1. ✗ Creatinine
2. ✗ Bilirubin
3. ✗ INR
4. ✓ Albumin

Question Number : 102 Question Id : 57996510054 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

CD 117 immunohistochemistry stain is used for

Options :

1. ✗ Melanoma
2. ✗ Angiosarcoma
3. ✓ Gastrointestinal stromal tumour
4. ✗ Neuroendocrine tumour

Question Number : 103 Question Id : 57996510055 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The most common benign tumor of the lung is

Options :

1. ✗ Adenoma
2. ✓ Hemartoma
3. ✗ Teratoma
4. ✗ Fibroma

Question Number : 104 Question Id : 57996510056 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Abbe- estlander flap is used for

Options :

1. ✘ Tongue
2. ✘ Eyelid
3. ✔ Lip
4. ✘ Ears

Question Number : 105 Question Id : 57996510057 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

First bite syndrome can occur as a complication of which of the following surgery?

Options :

1. ✔ Resection of carotid body tumour
2. ✘ Hemi-thyroidectomy
3. ✘ Surgery of soft palate
4. ✘ Surgery of pinna

Question Number : 106 Question Id : 57996510058 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common complication after splenectomy is

Options :

1. ✘ OPSI
2. ✘ Avascular necrosis of greater curvature of stomach'
3. ✘ Pancreatitis
4. ✔ Atelectasis

Question Number : 107 Question Id : 57996510059 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common type of thyroid carcinoma is

Options :

1. ✔ Papillary
2. ✘ Follicular
3. ✘ Medullary
4. ✘ Anaplastic

Question Number : 108 Question Id : 57996510060 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common malignancy of liver is

Options :

1. ✘ Hepatocellular carcinoma
2. ✘ Hepatoblastoma
3. ✘ Hemangioendothelioma
4. ✔ None of the given options are correct

Question Number : 109 Question Id : 57996510061 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Non correctable lesion after ulcerative colitis is

Options :

1. ✘ Skin lesions
2. ✘ Arthritis
3. ✔ Primary sclerosing cholangitis
4. ✘ Iritis

Question Number : 110 Question Id : 57996510062 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Stereotactic surgery is used for treatment of

Options :

1. ✔ Brain tumour
2. ✘ Lung carcinoma
3. ✘ Cervix cancer
4. ✘ Renal carcinoma

Question Number : 111 Question Id : 57996510063 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

What is the frequency of vibration of harmonic scalpel?

Options :

1. ✘ 2000-5000 Hz
2. ✘ 10000-15000 Hz
3. ✔ 20000-50000 Hz
4. ✘ 35000-70000 Hz

Question Number : 112 Question Id : 57996510064 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Causative organism for Asiatic cholangiohepatitis is

Options :

1. ✘ Brugia timori
2. ✔ Clonorchis sinensis
3. ✘ Ascaris lumbricoides
4. ✘ Echinococcus multilocularis

Question Number : 113 Question Id : 57996510065 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is true about acute graft rejection?

Options :

1. ✘ Caused by preformed antibody
2. ✘ It is not reversible
3. ✘ Vascular myointimal proliferation is seen
4. ✔ Predominantly T cell dependent

Question Number : 114 Question Id : 57996510066 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

T staging of esophageal carcinoma is best done by



**Options :**

1. ✓ EUS
2. ✗ MRI
3. ✗ CT
4. ✗ PET

**Question Number : 115 Question Id : 57996510067 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical**

**Correct : 1 Wrong : 0.33**

Most common site for lentigo melanoma is

**Options :**

1. ✗ Soles
2. ✓ Face
3. ✗ Legs
4. ✗ Trunk

**Question Number : 116 Question Id : 57996510068 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical**

**Correct : 1 Wrong : 0.33**

Jaboulay's operation is done for

**Options :**

1. ✗ Rectal prolapsed
2. ✗ Fistula in ano
3. ✓ Hydrocele
4. ✗ Inguinal hernia

**Question Number : 117 Question Id : 57996510069 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical**

**Correct : 1 Wrong : 0.33**

Which of the following is not true about torsion of testis?

**Options :**

1. ✗ Presents with sudden pain in testis
2. ✓ Commonly associated with pyuria
3. ✗ Doppler U/S shows decreased blood flow to testis
4. ✗ Simultaneous surgical exploration of the opposite testis is also to be done.

**Question Number : 118 Question Id : 57996510070 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical**

**Correct : 1 Wrong : 0.33**

What is the treatment of choice for adenocarcinoma of appendix measuring <1 cm in greatest diameter, not involving the base, without any lymphatic spread and peritoneal dissemination?

**Options :**

1. ✗ Appenicectomy only
2. ✗ Appendicectomy and post operative chemotherapy
3. ✗ Chemoradiation only
4. ✓ Right hemicolectomy

**Question Number : 119 Question Id : 57996510071 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical**

**Correct : 1 Wrong : 0.33**

Most common position of appendix is

Options :

1. ✘ Pelvic
2. ✘ Ileal
3. ✘ Subcaecal
4. ✔ Retrocaecal

Question Number : 120 Question Id : 57996510072 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Paget-Schroetter disease is caused by

Options :

1. ✘ Excessive breakdown and formation of bone, followed by disorganised bone remodelling.
2. ✔ upper extremity deep vein thrombosis
3. ✘ noninvasive intraepithelial (in the skin) adenocarcinoma outside the mammary gland
4. ✘ bilateral breast carcinoma involving nipple areola complex

Question Number : 121 Question Id : 57996510073 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Fogarty's catheter is used for

Options :

1. ✘ Suprapubic drainage of urinary bladder
2. ✘ Parenteral hyperalimentation
3. ✔ Removal of embolus from blood vessels
4. ✘ Ureteric catheterization

Question Number : 122 Question Id : 57996510074 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

What is the Ankle brachial index for imminent necrosis?

Options :

1. ✔  $< 0.3$
2. ✘  $< 0.5$
3. ✘  $< 0.9$
4. ✘  $< 1.0$

Question Number : 123 Question Id : 57996510075 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Fowler Stephen surgery is done for

Options :

1. ✘ Epispadias
2. ✘ Hypospadias
3. ✘ Exotrophy of bladder
4. ✔ Cryptorchodism

Question Number : 124 Question Id : 57996510076 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Radioactive iodine uptake is not seen in

Options :

1. ✘ Solitary toxic nodule
2. ✘ Follicular carcinoma thyroid
3. ✘ Papillary carcinoma of thyroid
4. ✔ Medullary carcinoma of thyroid

Question Number : 125 Question Id : 57996510077 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Migratory necrolytic skin rash is seen in?

Options :

1. ✘ ACTHoma
2. ✘ VIPoma
3. ✔ Glucagonoma
4. ✘ Somatostatinoma

Question Number : 126 Question Id : 57996510078 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The commonest cause of abdominal aortic aneurysm is?

Options :

1. ✘ Trauma
2. ✘ Syphilis
3. ✘ Vasculitis
4. ✔ Atherosclerosis

Question Number : 127 Question Id : 57996510079 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

First autologous renal transplantation was performed by?

Options :

1. ✔ Hardy
2. ✘ Higgins
3. ✘ Studor
4. ✘ Kavosis

Question Number : 128 Question Id : 57996510080 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is the most troublesome source of bleeding during a radical retropubic prostatectomy?

Options :

1. ✔ Dorsal venous complex
2. ✘ Inferior vesical pedicle
3. ✘ Superior vesical pedicle
4. ✘ Seminal vesicular artery

Question Number : 129 Question Id : 57996510081 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following catheter materials is most suited for long-term use?

Options :

1. ✘ Latex
2. ✔ Silicone
3. ✘ Rubber
4. ✘ Polyurethane

Question Number : 130 Question Id : 57996510082 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In which one of the following head and neck cancer, perineural invasion is most commonly seen?

Options :

1. ✘ Adenocarcinoma
2. ✔ Adenoid cystic carcinoma
3. ✘ Basal Cell Adenoma
4. ✘ Squamous cell carcinoma

Question Number : 131 Question Id : 57996510083 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common type of gall bladder stone is

Options :

1. ✘ Mixed
2. ✔ Cholesterol
3. ✘ Black pigmented
4. ✘ Brown pigmented

Question Number : 132 Question Id : 57996510084 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which is the first investigation for solitary thyroid nodule?

Options :

1. ✘ FNAC
2. ✘ USG- neck
3. ✔ Thyroid function test
4. ✘ Radioactive Iodine Scan

Question Number : 133 Question Id : 57996510085 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Portal hypertension is defined as portal venous pressure

Options :

1. ✘ > 5 mm Hg
2. ✔ >10 mm Hg
3. ✘ >15 mm Hg
4. ✘ >20 mm Hg

Question Number : 134 Question Id : 57996510086 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Tumour marker for Ca GB is

Options :

1. ✓ CA 19-9
2. ✗ CA- 125
3. ✗ CA - 72-4
4. ✗ B-HCG

Question Number : 135 Question Id : 57996510087 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Thyroid lymphoma is a risk factor for

Options :

1. ✗ Riedel's thyroiditis
2. ✓ Hashimoto's thyroiditis
3. ✗ Subacute thyroiditis
4. ✗ Acute thyroiditis

Question Number : 136 Question Id : 57996510088 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All are done in whipples operation except

Options :

1. ✗ Pancreaticojejunostomy
2. ✗ Hepaticojejunostomy
3. ✗ Gastrojejunostomy
4. ✓ Duodenojejunostomy

Question Number : 137 Question Id : 57996510089 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Widening of C- Loop of duodenum in x-ray is diagnostic of

Options :

1. ✗ Calculus in ampulla of water
2. ✓ Carcinoma head of the pancreas
3. ✗ Chronic pancreatitis
4. ✗ Periampullary carcinoma

Question Number : 138 Question Id : 57996510090 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Nelson syndrome is most commonly seen after

Options :

1. ✗ Hypophysectomy
2. ✓ Adrenalectomy
3. ✗ Thyroidectomy
4. ✗ Orchidectomy

Question Number : 139 Question Id : 57996510091 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following does not occur with gastrinoma?

Options :

1. ✘ Epigastric pain
2. ✘ Diarrhea
3. ✔ Basal acid output <15 mEq/L
4. ✘ Serum gastrin > 200 pg/ml

Question Number : 140 Question Id : 57996510092 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Meckel's diverticulum is remnant of

Options :

1. ✘ Stenson duct
2. ✘ Wolffian duct
3. ✘ Mullerian duct
4. ✔ Vitelointestinal duct

Question Number : 141 Question Id : 57996510093 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common site for parathyroid adenoma is

Options :

1. ✘ Intra thyroid
2. ✘ In the superior parathyroid lobe
3. ✔ In the inferior parathyroid lobe
4. ✘ Mediastinal

Question Number : 142 Question Id : 57996510094 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A Warthin's tumour is

Options :

1. ✔ An adenolymphoma of parotid gland
2. ✘ A pleomorphic adenoma of the parotid
3. ✘ A carcinoma of the parotid gland
4. ✘ A carcinoma of the submandibular salivary gland

Question Number : 143 Question Id : 57996510095 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All of the following constitute the Saint's triad except

Options :

1. ✘ Gall stones
2. ✔ Jaundice
3. ✘ Hiatus hernia
4. ✘ Colonic diverticulosis

Question Number : 144 Question Id : 57996510096 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Zollinger Ellison syndrome is characterized by all except

Options :

1. ✘ Profound gastric hypersecretion
2. ✘ Diarrhea with occasional steatorrhea
3. ✔ Hypocalcemia
4. ✘ Hypergastrinemia

Question Number : 145 Question Id : 57996510097 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Subclavian steal syndrome is due to

Options :

1. ✘ Occlusion/stenosis of the carotid artery
2. ✘ Occlusion/stenosis of the vertebral artery
3. ✔ Occlusion of subclavian artery proximal to origin of Vertebral artery
4. ✘ Occlusion of subclavian artery distal to origin of Vertebral artery

Question Number : 146 Question Id : 57996510098 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Regarding laparoscopic cholecystectomy which of the following is correct?

Options :

1. ✘ It is primarily done for cholecystitis in 3rd trimester of pregnancy
2. ✔ It has higher incidence of bile duct injury than open cholecystectomy
3. ✘ It is contraindicated in acute cholecystitis
4. ✘ It is safer than open cholecystectomy in patients with cardiorespiratory distress

Question Number : 147 Question Id : 57996510099 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Horner's syndrome include all except

Options :

1. ✘ Miosis
2. ✘ Anhydrosis
3. ✘ Hyperchromatic iris
4. ✔ Apparent Exophthalmos

Question Number : 148 Question Id : 57996510100 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common type of shock in emergency room is

Options :

1. ✘ Cardiogenic
2. ✔ Hypovolemic
3. ✘ Neurogenic
4. ✘ Septic

Question Number : 149 Question Id : 57996510101 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In 3rd degree burn all are seen except

**Options :**

1. ✘ Vesicles are absent
2. ✔ Painful
3. ✘ Leathery skin
4. ✘ Reddish due to Hb infiltration

Question Number : 150 Question Id : 57996510102 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes  
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Bisgard treatment is for

**Options :**

1. ✘ Arterial ulcer
2. ✔ Venous ulcer
3. ✘ TAO
4. ✘ Raynaud's phenomenon