

Question Paper Preview

Notations :

- 1.Options shown in green color and with ✓ icon are correct.
- 2.Options shown in red color and with ✗ icon are incorrect.

Question Paper Name:	Civil Assistant Surgeon in Gynecology 5th Dec 2017 Shift 1
Subject Name:	Civil Assistant Surgeon in Gynecology
Creation Date:	2017-12-05 14:04:29
Duration:	150
Total Marks:	150
Display Marks:	No
Calculator:	Scientific
Magnifying Glass Required?:	No
Ruler Required?:	No
Eraser Required?:	No
Scratch Pad Required?:	No
Rough Sketch/Notepad Required?:	No
Protractor Required?:	No

Civil Assistant Surgeon in Gynecology

Group Number :	1
Group Id :	57996568
Group Maximum Duration :	0
Group Minimum Duration :	150
Revisit allowed for view? :	No
Revisit allowed for edit? :	No
Break time:	0
Group Marks:	150

Civil Assistant Surgeon in Gynecology

Section Id :	57996568
Section Number :	1
Section type :	Online
Mandatory or Optional:	Mandatory
Number of Questions:	150
Number of Questions to be attempted:	150
Section Marks:	150
Display Number Panel:	Yes
Group All Questions:	No

Sub-Section Number:	1
Sub-Section Id:	579965106
Question Shuffling Allowed :	Yes

Question Number : 1 Question Id : 57996510103 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is responsible for pubertal growth in female?

Options :

1. ✘ Decreased level of adrenal androgens at puberty
2. ✘ High level of estrogen at puberty
3. ✔ Pulsatile release of GnRH during sleep
4. ✘ Increased sensitivity of HPO axis to estrogen

Question Number : 2 Question Id : 57996510104 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Medication used in treatment of idiopathic central precocious puberty is:

Options :

1. ✘ Exogenous gonadotrophins
2. ✘ Levonor gestrel
3. ✔ Gn RH analogues
4. ✘ Ethinyl estradiol

Question Number : 3 Question Id : 57996510105 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

pH of vagina during reproductive life is

Options :

1. ✘ 3.0 - 3.5
2. ✔ 4.0-4.5
3. ✘ 6.0-7.0
4. ✘ > 7.0

Question Number : 4 Question Id : 57996510106 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

17 year-old girl with amenorrhea, atrophied breast hypo plastic uterus

Options :

1. ✔ Turner's syndrome
2. ✘ Gonadal dysgenesis
3. ✘ Androgen insensitivity syndrome
4. ✘ Klinefelter's syndrome

Question Number : 5 Question Id : 57996510107 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Pure gonadal dysgenesis will be diagnosed in the presence of:

Options :

1. ✔ Bilateral Streak gonads
2. ✘ Bilateral dysgenetic gonads
3. ✘ One side steak and other digenetic gonads
4. ✘ One side steak and other normal looking gonad

Question Number : 6 Question Id : 57996510108 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Precocious puberty may be seen in all of the following condition except:

Options :

1. ✘ Granulosa-cell tumour
2. ✘ Head-injury
3. ✘ Corticosteroid intake
4. ✔ Hyperthyroidism

Question Number : 7 Question Id : 57996510109 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A 25- year old female with history of multiple contacts presenting with growth on vulva, the probable diagnosis is:

Options :

1. ✔ Condyloma accuminata
2. ✘ Veeruca plana
3. ✘ Veeruca vulgaris
4. ✘ Condyloma lata

Question Number : 8 Question Id : 57996510110 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All are clinical features of PID except

Options :

1. ✘ Temp > 38 C
2. ✘ WBC count of 15,000
3. ✔ ESR - 10 mm/hour
4. ✘ Tenderness on movement of cervix

Question Number : 9 Question Id : 57996510111 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common route of transmission of endometrial tuberculosis is:

Options :

1. ✘ Direct local spread
2. ✘ Lymphatic spread
3. ✘ Retrograde spread
4. ✔ Hematogenous spread

Question Number : 10 Question Id : 57996510112 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A lady with prolapsed uterus after fothergill's repair will complain of following except:

Options :

1. ✔ First trimester abortion
2. ✘ Cervical dystocia
3. ✘ Premature labour
4. ✘ Premature rupture of membrane

Question Number : 11 Question Id : 57996510113 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common site of obstetric injury leading to uretero vaginal fistula.

Options :

1. ✘ Infundibulo pelvic ligament
2. ✘ Vaginal vault
3. ✘ Ureteric tunnel
4. ✔ Below cardinal ligament where uterine artery crosses

Question Number : 12 Question Id : 57996510114 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common genital prolapse is:

Options :

1. ✔ Cystocele
2. ✘ Procidentia
3. ✘ Rectocele
4. ✘ Enterocoele

Question Number : 13 Question Id : 57996510115 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Shirodkar sling operation may be associated with all complications except:

Options :

1. ✔ Enterocoele
2. ✘ Subacute intestinal obstructions
3. ✘ Clreteral injury
4. ✘ Parasthesia over inner aspect.

Question Number : 14 Question Id : 57996510116 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Corkscrew shaped endometrial glands are seen in

Options :

1. ✘ Early proliferative phase
2. ✘ Late proliferative phase
3. ✘ Early secretory phase
4. ✔ Late secretory phase

Question Number : 15 Question Id : 57996510117 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Best management of vault prolapse is:

Options :

1. ✔ Sacral colpopexy
2. ✘ Sacrospinous ligament fixation
3. ✘ Le forts repair
4. ✘ Anterior colporapphy

Question Number : 16 Question Id : 57996510118 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following surgeries for stress incontinence has highest success rate

Options :

1. ✘ Burch colposuspension
2. ✘ Pereyra sling
3. ✘ Kelly's stitch
4. ✔ Tension free vaginal tape (TVT)

Question Number : 17 Question Id : 57996510119 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The risk of asherman syndrome is the highest if dilation and curettage (D and C) is done for the following condition

Options :

1. ✘ Medical termination of pregnancy
2. ✘ Missed abortion
3. ✘ Dysfunctional uterine bleeding
4. ✔ Post partum hemorrhage

Question Number : 18 Question Id : 57996510120 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Fallopian tube dysmotility is seen in

Options :

1. ✘ Noonan syndrome
2. ✘ Turner syndrome
3. ✔ Kartagener syndrome
4. ✘ Marfan syndrome

Question Number : 19 Question Id : 57996510121 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Post coital test showing non motile sperms in the cervical smear and motile sperms from the posterior fornix suggests:

Options :

1. ✘ Faulty coital Practice
2. ✔ Immunological defect
3. ✘ Hypospadias
4. ✘ Azoospermia

Question Number : 20 Question Id : 57996510122 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which is a not an essential criteria according to WHO for normal semen analysis:

Options :

1. ✘ Sperm count > 20 million/ml
2. ✔ Volume > 1ml
3. ✘ Sperm with normal morphology (strict criteria) > 15%

4. ✘ Motility > 25% with rapidly progressive motility

Question Number : 21 Question Id : 57996510123 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Spermicidal jelly acts through

Options :

1. ✘ Acrosomal enzyme
2. ✘ Cervical Enzyme alteration
3. ✘ Glucose uptake inhibition by sperms
4. ✔ Disruption of cell membrane

Question Number : 22 Question Id : 57996510124 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Muscle that is not inserted into the perineal body is

Options :

1. ✘ Levator Ani
2. ✘ Ischiocavernosus
3. ✘ Bulbospongiosus
4. ✔ Superficial perineal muscles

Question Number : 23 Question Id : 57996510125 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A Patient had a spontaneous abortion, then she came with amenorrhea and FSH 6 IU/ ml what the most probably diagnosis?

Options :

1. ✘ Ovarian failure
2. ✔ Uterine synechiae
3. ✘ Pregnancy
4. ✘ Pituitary failure

Question Number : 24 Question Id : 57996510126 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The most common cause of secondary amenorrhea in India is:

Options :

1. ✔ Endometrial tuberculosis
2. ✘ Premature ovarian failure
3. ✘ Polycystic ovarian syndrome
4. ✘ Sheehan's syndrome

Question Number : 25 Question Id : 57996510127 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In DUB, there is:

Options :

1. ✔ Increased estrogen
2. ✘ Decreased receptors of progesterone

- ✘ Decreased receptors of estrogen
- ✘ Pituitary imbalance of hormones

Question Number : 26 Question Id : 57996510128 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common endometrial pattern in dysfunctional uterine bleeding:

Options :

- ✔ Normal
- ✘ Hyperplastic with swiss-cheese pattern
- ✘ Nonsecretory
- ✘ Atrophic

Question Number : 27 Question Id : 57996510129 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Primary amenorrhoea is most commonly associated with

Options :

- ✘ Developmental defect of the genital tract
- ✘ Tuberculosis
- ✘ Endocrine disorders
- ✔ Chromosomal Abnormality

Question Number : 28 Question Id : 57996510130 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

To diagnose uterus didelphys, procedure of choice is :

Options :

- ✘ Laparoscopy
- ✘ IVP
- ✘ HSG
- ✔ USG

Question Number : 29 Question Id : 57996510131 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Ideal age for repair of vaginal agenesis is:

Options :

- ✘ 6 months
- ✘ 3 year
- ✘ At puberty
- ✔ Before marriage

Question Number : 30 Question Id : 57996510132 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All of the following are features of mullerian agenesis except:

Options :

- ✘ 46XX karyotype
- ✘ Normal breast development

- ✘ Absent vagina
- ✔ Ovarian agenesis

Question Number : 31 Question Id : 57996510133 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Beaded appearance of fallopian tube is seen in

Options :

- ✘ Carcinoma of tubes
- ✘ Chlamydial infection
- ✘ Gonorrhoea
- ✔ Tuberculosis

Question Number : 32 Question Id : 57996510134 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Mayer Rokitansky Kuster Hauser syndrome consists of:

Options :

- ✘ Ovaries, Uterus fallopian tubes present
- ✔ Uterus absent, Fallopian tube ovaries present
- ✘ All absent
- ✘ Uterus present tubes and ovaries absent

Question Number : 33 Question Id : 57996510135 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following pubertal events in girls is not estrogen dependent:

Options :

- ✘ Menstruation
- ✘ Vaginal Cornification
- ✘ Height Spurt
- ✔ Hair Growth

Question Number : 34 Question Id : 57996510136 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common cause of ambiguous genitalia in female child is:

Options :

- ✘ Placenta steroid sulphatase deficiency
- ✘ Fetal aromatase deficiency
- ✘ Wnt4 mutation
- ✔ Congenital adrenal hyperplasia

Question Number : 35 Question Id : 57996510137 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Best prenatal treatment for CAH is:

Options :

- ✔ Dexamethasone
- ✘ Betamethasone

3. ✘ Prednisolone
4. ✘ Hydrocortisone

Question Number : 36 Question Id : 57996510138 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Ring pessary for prolapse is indicated in

Options :

1. ✘ Permanent cure
2. ✘ Young patients
3. ✘ Cure for dyspareunia
4. ✔ Puerperium

Question Number : 37 Question Id : 57996510139 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Young male presents with delayed puberty with decreased FSH, LH and testosterone. Which of the following is NOT possible?

Options :

1. ✘ Kallmann Syndrome
2. ✔ Klinefelter's syndrome
3. ✘ Constitutional delay
4. ✘ Dax-1 gene mutation.

Question Number : 38 Question Id : 57996510140 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common fistula as a complication of obstructed labour is

Options :

1. ✔ Vesico Vaginal fistula
2. ✘ Utero Vaginal fistula
3. ✘ Uretero Abdominal fistula
4. ✘ Vesico Uterine fistula

Question Number : 39 Question Id : 57996510141 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Uterine Artery embolization is done using

Options :

1. ✘ Thrombin
2. ✔ Polyvinyl alcohol
3. ✘ Vitamin K
4. ✘ Iodine

Question Number : 40 Question Id : 57996510142 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Use of OCP's are known to protect against following malignancies except:

Options :

1. ✘ Ovarian carcinoma

- ✘ Endometrial carcinoma
- ✘ Uterine sarcoma
- ✔ Carcinoma cervix

Question Number : 41 Question Id : 57996510143 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True statement about minipill is:

Options :

- ✔ Irregular Vaginal Bleeding may be a side effect.
- ✘ Used in with combination with oral contraceptive pills
- ✘ Cannot be used in lactation
- ✘ Prevents ectopic pregnancy

Question Number : 42 Question Id : 57996510144 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Composition of Nova-T:

Options :

- ✔ Copper and Silver
- ✘ Copper and aluminium
- ✘ Copper only
- ✘ Copper and Selenium.

Question Number : 43 Question Id : 57996510145 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Contraceptive to be avoided in epilepsy:

Options :

- ✔ OCP
- ✘ Condom
- ✘ IUCD
- ✘ Mirena

Question Number : 44 Question Id : 57996510146 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The intra-abdominal pressure during laparoscopy should be set between:

Options :

- ✘ 5-8 mm hg
- ✔ 10-15 mm hg
- ✘ 20-25 mm hg
- ✘ 30-35 mm hg

Question Number : 45 Question Id : 57996510147 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Least failure in sterilization occurs with

Options :

- ✘ Falope ring

- ✘ Bipolar cautery
- ✔ Unipolar cauterisation
- ✘ Hulka clip

Question Number : 46 Question Id : 57996510148 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Not true about red degeneration of myomas is:

Options :

- ✘ It occurs commonly during pregnancy
- ✔ Immediate surgical intervention is needed
- ✘ Due to interference with blood supply
- ✘ Treated with analgesics

Question Number : 47 Question Id : 57996510149 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

To start with all Fibroids are :

Options :

- ✔ Interstitial
- ✘ Submucous
- ✘ Subserous
- ✘ Ovarian

Question Number : 48 Question Id : 57996510150 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Uterine fibromyoma is associated with:

Options :

- ✔ Endometriosis
- ✘ Pelvic inflammatory disease.
- ✘ Ovarian Ca
- ✘ Amenorrhea

Question Number : 49 Question Id : 57996510151 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Investigation of choice in post-menopausal bleeding is

Options :

- ✘ D&C
- ✘ Colposcopy
- ✘ PAP Smear
- ✔ Fractional Curettage with Cervix Biopsy

Question Number : 50 Question Id : 57996510152 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Emergency contraception mechanisms include all except

Options :

- ✔ Inhibits ovulation

2. ✘ Prevents implantation
3. ✘ Interferes with early pregnancy
4. ✘ Inhibits fertilisation

Question Number : 51 Question Id : 57996510153 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Pain in endometriosis correlates with:

Options :

1. ✔ A depth of Invasion
2. ✘ Multiple Sites
3. ✘ CA 125
4. ✘ A stage of disease

Question Number : 52 Question Id : 57996510154 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Peg cells are seen in

Options :

1. ✘ Endometrium
2. ✘ Cervix
3. ✔ Fallopian tubes
4. ✘ Ovary

Question Number : 53 Question Id : 57996510155 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True regarding adenomyosis is:

Options :

1. ✘ Most common is nullipara
2. ✘ Progestin are agents of choice for medical management
3. ✔ Presents with menorrhagia dysmenorrhoea and an enlarged uterus.
4. ✘ More common in young women

Question Number : 54 Question Id : 57996510156 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following modalities have shown best result for pre menstrual syndrome.?

Options :

1. ✔ SSRI
2. ✘ Progesterone
3. ✘ Oestrogen
4. ✘ Anxiolytics

Question Number : 55 Question Id : 57996510157 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The commonest cause of primary amenorrhoea is:

Options :

1. ✘ Genital Tuberculosis

2. ✓ Ovarian Dysgenesis
3. ✗ Mullerian duct anomalies
4. ✗ Hypothyroidism

Question Number : 56 Question Id : 57996510158 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which is not primary amenorrhoea?

Options :

1. ✓ Sheehan's syndrome.
2. ✗ Kallmann's syndrome.
3. ✗ Mayer Rokitansky koster hauser syndrome.
4. ✗ Turner syndrome.

Question Number : 57 Question Id : 57996510159 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

"Today" sponge contains

Options :

1. ✓ Nonoxynol
2. ✗ Delfen
3. ✗ Femshield
4. ✗ Desogestrel

Question Number : 58 Question Id : 57996510160 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Commonest cause of post menopausal bleeding in india is:

Options :

1. ✗ Ca endometrium
2. ✓ Ca cervix
3. ✗ Ca vulva
4. ✗ Ovarian tumor.

Question Number : 59 Question Id : 57996510161 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A teenage girl presented in OPD with moderate acne and history of irregular menses. What treatment will you give ?

Options :

1. ✗ Oral isotretinoin.
2. ✗ Oral acitretin.
3. ✗ Oral minocycline.
4. ✓ cyproterone acetate.

Question Number : 60 Question Id : 57996510162 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Lady recovered from severe PPH, complains of failure of lactation and menstruation, Which of the following can be seen :

Options :

1. ✓ Increased excretion of Na⁺
2. ✗ Retention of water.
3. ✗ Increased Prolactin.
4. ✗ Increased GnRH.

Question Number : 61 Question Id : 57996510163 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The risk of endometrial cancer is highest with the following histological pattern of endometrial hyperplasia :

Options :

1. ✗ Simple hyperplasia without atypia.
2. ✗ Simple hyperplasia with atypia.
3. ✗ Complex hyperplasia without atypia.
4. ✓ Complex hyperplasia with atypia.

Question Number : 62 Question Id : 57996510164 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Long term tamoxifen therapy may cause:

Options :

1. ✓ Endometrial Ca
2. ✗ Ovary Ca
3. ✗ Cervix Ca
4. ✗ Vagina Ca.

Question Number : 63 Question Id : 57996510165 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Point B in the treatment of carcinoma cervix receives the dose of:

Options :

1. ✗ 7000 cGy
2. ✓ 6000 cGy
3. ✗ 5000 cGy
4. ✗ 10000 cGy

Question Number : 64 Question Id : 57996510166 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Treatment of Stage III B carcinoma cervix is:

Options :

1. ✗ Wertheim procedure.
2. ✗ Schauta's procedure.
3. ✗ Chemotherapy.
4. ✓ Intracavitary Brachy therapy followed by external beam RT.

Question Number : 65 Question Id : 57996510167 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All of the following changes are seen in dysplasia a of squamocolumnar junction , except:

Options :

1. ✓ Breaking of basement membrane.
2. ✗ Change of epithelium.
3. ✗ Hyperchromatic nuclei.
4. ✗ Increased mitotic figure.

Question Number : 66 Question Id : 57996510168 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In a cervix low grade squamous intraepithelial lesion(LSIL) in bethseda system includes :

Options :

1. ✓ CIN I
2. ✗ CIN II
3. ✗ CIN III
4. ✗ Squamous metaplasia.

Question Number : 67 Question Id : 57996510169 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Therapeutic conisation is indicated in :

Options :

1. ✓ Microinvasive carcinoma.
2. ✗ CIN (III)
3. ✗ Unsatisfactory colposcopy withcervical dysplasia.
4. ✗ Cervical metaplasia.

Question Number : 68 Question Id : 57996510170 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is not a risk factor for CA cervix?

Options :

1. ✓ Low parity.
2. ✗ Multiple sexual partner.
3. ✗ Early sexual intercourse (<16 years)
4. ✗ Smoking.

Question Number : 69 Question Id : 57996510171 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common Ovarian tumor in less than 20 years is :

Options :

1. ✗ Epithelial tumor.
2. ✓ Germ cell tumor.
3. ✗ Metastatic tumor.
4. ✗ Sexcordstromal tumor.

Question Number : 70 Question Id : 57996510172 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following are masculinising tumors of the ovary :

Options :

1. ✘ Granulosa Cell tumor.
2. ✘ Dysgerminoma.
3. ✘ Dermoid Cyst.
4. ✔ Arrhenoblastoma.

Question Number : 71 Question Id : 57996510173 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All the components of Meig's Syndrome, except:

Options :

1. ✘ Pleural effusion.
2. ✘ Ovarian Tumor.
3. ✘ Ascites.
4. ✔ Pericardial Effusion.

Question Number : 72 Question Id : 57996510174 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common Vaginal carcinoma is :

Options :

1. ✔ Squamous Cell Carcinoma.
2. ✘ Adenocarcinoma.
3. ✘ Botroid tumor.
4. ✘ Columnar hyperplasia.

Question Number : 73 Question Id : 57996510175 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Common differential diagnosis of verrucous carcinoma is :

Options :

1. ✘ condylomata lata.
2. ✔ Condylomata accuminata.
3. ✘ Adenocarcinoma.
4. ✘ Tuberculosis

Question Number : 74 Question Id : 57996510176 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Pyometra is a complication associated with all of the following conditions, except:

Options :

1. ✔ Carcinoma vulva.
2. ✘ Carcinoma Cervix.
3. ✘ Carcinoma endometrium.
4. ✘ pelvic radiotherapy.

Question Number : 75 Question Id : 57996510177 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Cornification index or eosinophilic index indicates :

Options :

1. ✘ Progesterone effect
2. ✔ Estrogenic effect
3. ✘ Effect of LH
4. ✘ All of the given options are correct

Question Number : 76 Question Id : 57996510178 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following is the least likely physiological change in pregnancy:

Options :

1. ✘ Increase in intravascular volume.
2. ✘ Increase in cardiac output.
3. ✘ Increase in stroke volume.
4. ✔ Increase in peripheral vascular resistance.

Question Number : 77 Question Id : 57996510179 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which cardiovascular change is physiological in last trimester of pregnancy:

Options :

1. ✘ Mid diastolic murmur.
2. ✘ Occasional atrial fibrillation.
3. ✔ Shift of apical impulse laterally and upwards in the left 4th intercostal space.
4. ✘ Cardiomegaly.

Question Number : 78 Question Id : 57996510180 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All of the following may be observed in normal pregnancy except:

Options :

1. ✘ Fall in serum iron concentration.
2. ✘ Increase in serum iron binding capacity.
3. ✔ Increase in blood viscosity.
4. ✘ Increase in blood oxygen carrying capacity.

Question Number : 79 Question Id : 57996510181 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

During fetal life maximum growth is caused by:

Options :

1. ✘ Growth hormone.
2. ✔ Insulin.
3. ✘ Cortisol.
4. ✘ Thyroxin

Question Number : 80 Question Id : 57996510182 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Best test for estimating hCG:

Options :

1. ✓ Radio Immunoassay.
2. ✗ ELISA.
3. ✗ Radio receptor assay.
4. ✗ Bioassay.

Question Number : 81 Question Id : 57996510183 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The clotting factor which is not increased in pregnancy:

Options :

1. ✗ Factor 2.
2. ✗ Factor 7.
3. ✗ Factor 10.
4. ✓ Factor 11.

Question Number : 82 Question Id : 57996510184 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A prosthetic valve patient switches to heparin at what time during pregnancy:

Options :

1. ✗ 28 wks.
2. ✗ 32 wks.
3. ✓ 36 wks.
4. ✗ Postpartum.

Question Number : 83 Question Id : 57996510185 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The role of placental lactogen is:

Options :

1. ✗ Stimulate milk production.
2. ✗ Fetal breast development.
3. ✗ Growth of fetus.
4. ✓ Endocrine regulation.

Question Number : 84 Question Id : 57996510186 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The term "placental sign" denotes:

Options :

1. ✗ Alteration of FHR on pressing head into the pelvis.
2. ✓ Spotting on the expected date of period in early months of pregnancy.
3. ✗ Permanent lengthening of the cord in 3rd stage of labour.
4. ✗ Slight gush of bleeding on the 3rd stage of labour.

Question Number : 85 Question Id : 57996510187 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In transvaginal ultrasound, earliest detection of gestation sac is by:

Options :

- ✘ 21 days after ovulation.
- ✘ 21 days after implantation.
- ✘ 28 days post ovulation.
- ✔ 14 days after ovulation.

Question Number : 86 Question Id : 57996510188 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Kegels exercise should begin: Version 1:

Options :

- ✘ Immediately after delivery.
- ✔ 24 hrs after delivery.
- ✘ 3 weeks after delivery.
- ✘ 6 weeks after delivery.

Question Number : 87 Question Id : 57996510189 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Quickening is first observed in a primi by

Options :

- ✔ 18 weeks
- ✘ 16 weeks
- ✘ 30 weeks
- ✘ 10 weeks

Question Number : 88 Question Id : 57996510190 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Minimum level of beta HCG detected by radioimmunoassay is:

Options :

- ✘ 1.5 - 3.5 IU/ml
- ✘ 0.5 - 1.0 IU/ml
- ✘ 0.02 - 0.05 IU/ml
- ✔ 0.001 IU/ml

Question Number : 89 Question Id : 57996510191 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Sensitivity of uterine musculature:

Options :

- ✘ Enhanced by progesterone.
- ✘ Enhanced by estrogen.
- ✘ Inhibited by estrogen.
- ✔ Enhanced by estrogen and inhibited by progesterone.

Question Number : 90 Question Id : 57996510192 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which is not included in the Active Management of III stage of labor?

Options :

- ✘ Uterotonic within 1 minute of delivery.
- ✔ Immediate clamping, cutting and ligation of cord.
- ✘ General massage of uterus.
- ✘ Controlled cord traction.

Question Number : 91 Question Id : 57996510193 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Pain in early labor is limited to dermatomes:

Options :

- ✔ T10-L1.
- ✘ S1-S3.
- ✘ L4-L5.
- ✘ L2-L3.

Question Number : 92 Question Id : 57996510194 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Ritgen maneuver is done in:

Options :

- ✘ Shoulder dystocia.
- ✘ For delivery of head in breech presentation.
- ✘ For delivery of legs in breech.
- ✔ For delivery of head in normal labour.

Question Number : 93 Question Id : 57996510195 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All are true about origin and propagation of contractions except:

Options :

- ✘ The right pacemaker predominates over left.
- ✔ Intensity of propagation is greatest at cervix.
- ✘ The contraction spreads from pacemaker towards cervix.
- ✘ Speed of contraction in 2cm/sec.

Question Number : 94 Question Id : 57996510196 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The pre requisites for internal rotation of the head are all except:

Options :

- ✘ Well-flexed head.
- ✘ Efficient uterine contraction.
- ✘ Favorable shape of the pelvis.
- ✔ Tone of the abdominal muscles.

Question Number : 95 Question Id : 57996510197 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Cervical incompetence is diagnosed if No.8 Hegar dilator can be passed beyond internal os without resistance

Options :

1. ✘ During 14th Week of pregnancy
2. ✘ During 6th week of pregnancy
3. ✔ In the pre-menstrual period
4. ✘ In the pre-ovulatory period

Question Number : 96 Question Id : 57996510198 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Single Umbilical Artery is associated with

Options :

1. ✔ Fetal congenital anomalies
2. ✘ Diabetes
3. ✘ Intra Uterine Infection
4. ✘ Hydrocephalus

Question Number : 97 Question Id : 57996510199 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All the following are used for induction of labor except:

Options :

1. ✔ PG F2 alpha tablet.
2. ✘ PG E1 tablet.
3. ✘ PG E2 gel.
4. ✘ Misoprostol.

Question Number : 98 Question Id : 57996510200 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Benefits of surgical induction (ARM) are all except:

Options :

1. ✘ Lowers the blood pressure in preeclampsia.
2. ✘ Relieves the maternal distress in hydramnios.
3. ✔ Decreases incidence of amnionitis
4. ✘ Reduces the need of caesarean section.

Question Number : 99 Question Id : 57996510201 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which one of the following methods for induction of labor should not be used in patient with previous lower segment caesarean section?

Options :

1. ✘ Prostaglandin gel
2. ✔ Prostaglandin tablet
3. ✘ Stripping of the membrane.
4. ✘ Oxytocin drip.

Question Number : 100 Question Id : 57996510202 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A lady with previous CS presents in labour. Trial of scar is contradicted in all except:

Options :

1. ✓ H/O previous LSCS due to malpresentation.
2. ✗ Knowing the fact that the previous CS was due to CPD.
3. ✗ H/O previous classical CS
4. ✗ Previous 3 LSCS

Question Number : 101 Question Id : 57996510203 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All of the following are true about augmentation of labour except:

Options :

1. ✓ twin pregnancy precludes the use of oxytocin.
2. ✗ amniotomy decreases the need for oxytocin use.
3. ✗ methods of augmentation doesn't increase the risk of operational management.
4. ✗ associated with risk of uterine hyperstimulation.

Question Number : 102 Question Id : 57996510204 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Lochia in correct order: during puerperium.

Options :

1. ✓ Rubra, Serosa, Alba.
2. ✗ Serosa, rubra, alba.
3. ✗ Alba, Serosa, rubra.
4. ✗ Alba, mucosa, serosa.

Question Number : 103 Question Id : 57996510205 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A pregnant female has past history of embolism in puerperium. What medical management she should take in next pregnancy to avoid this:

Options :

1. ✗ Compulsory prophylaxis with warfarin start at 10 weeks.
2. ✓ To take warfarin after delivery.
3. ✗ Chance of thromboembolism increases by 12% in the next pregnancy.
4. ✗ Does not need anything.

Question Number : 104 Question Id : 57996510206 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In comparison to breast milk, colostrum has higher content of:

Options :

1. ✗ Carbohydrates.
2. ✗ Fat.
3. ✓ Sodium
4. ✗ Potassium.

Question Number : 105 Question Id : 57996510207 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In ultrasound, abdominal circumference is measured at the level of

Options :

1. ✓ Stomach and Umbilical Vein
2. ✗ Kidney and Renal Artery
3. ✗ Spleen and Coeliac Plexus
4. ✗ Liver and Cystic Duct

Question Number : 106 Question Id : 57996510208 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All are complication of formula fed baby over human milk fed baby except:

Options :

1. ✗ Necrotizing enterocolitis
2. ✗ Otitis media
3. ✗ Hypokalemia.
4. ✓ Vit K deficiency.

Question Number : 107 Question Id : 57996510209 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Variable deceleration in CTG indicates

Options :

1. ✓ Cord compression
2. ✗ Head compression
3. ✗ Fetal Hypoxia
4. ✗ Fetal Death

Question Number : 108 Question Id : 57996510210 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In case of a recurrent spontaneous abortion, following investigation is unwanted:

Options :

1. ✗ Hysteroscopy.
2. ✗ Testing antiphospholipid antibodies.
3. ✓ Testing of TORCH infections.
4. ✗ Thyroid function tests.

Question Number : 109 Question Id : 57996510211 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A woman with H/O recurrent abortions present with isolated increase in APTT. Most likely cause is:

Options :

1. ✓ Lupus anticoagulant.
2. ✗ Factor VII
3. ✗ Von Willebrand's disease.
4. ✗ Hemophilia A.

Question Number : 110 Question Id : 57996510212 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

All of the following drugs have been used for medical abortion except:

Options :

1. ✘ Mifepristone.
2. ✘ Misoprostol.
3. ✘ Methotrexate.
4. ✔ Atosiban.

Question Number : 111 Question Id : 57996510213 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Blighted ovum is:

Options :

1. ✘ Synaptic Knobs.
2. ✔ Avascular villi.
3. ✘ Intervillous hemorrhage.
4. ✘ None of the given options are correct

Question Number : 112 Question Id : 57996510214 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Common cause of Secondary PPH is

Options :

1. ✘ Uterine atony
2. ✘ Uterine rupture
3. ✘ Vaginal laceration
4. ✔ Retained placental bits

Question Number : 113 Question Id : 57996510215 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In which part of fallopian tube ectopic pregnancy will have longest survival:

Options :

1. ✘ Isthmus.
2. ✘ Ampulla.
3. ✘ Cornua.
4. ✔ Interstitium.

Question Number : 114 Question Id : 57996510216 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Biochemical markers to detect aneuploidy in first trimester is

Options :

1. ✘ Relaxin
2. ✘ Inhibin A
3. ✔ PAPP - A
4. ✘ Alpha Feto Protein

Question Number : 115 Question Id : 57996510217 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Spiegelberg Criteria is used for diagnosis of

Options :

1. ✘ Ampullary pregnancy
2. ✘ Cervical pregnancy
3. ✔ Ovarian pregnancy
4. ✘ Intra uterine pregnancy

Question Number : 116 Question Id : 57996510218 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common infection resulting in ectopic pregnancy is

Options :

1. ✔ Chlamydia
2. ✘ Gonococci
3. ✘ Syphilis
4. ✘ Candida

Question Number : 117 Question Id : 57996510219 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Arias Stella reaction is not seen in:

Options :

1. ✘ Ovarian pregnancy,
2. ✘ Molar pregnancy,
3. ✘ Interstitial pregnancy.
4. ✔ Salpingitis isthmica nodosa.

Question Number : 118 Question Id : 57996510220 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

False about partial mole

Options :

1. ✘ caused by triploidy
2. ✘ can be diagnosed very early by USG
3. ✘ can be present as missed abortion
4. ✔ Rarely causes persistent GTD

Question Number : 119 Question Id : 57996510221 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A case of gestational trophoblastic neoplasia belongs to high risk group disease, if disease develops after

Options :

1. ✘ hydatidiform mole
2. ✔ Full term pregnancy
3. ✘ Spontaneous abortion
4. ✘ Ectopic pregnancy

Question Number : 120 Question Id : 57996510222 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Commonest manifestation in pre-term infant is

Options :

- ✘ Parchment skin
- ✘ Abundant scalp hair
- ✔ Lanugo hair
- ✘ Vallus hair

Question Number : 121 Question Id : 57996510223 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Choriocarcinoma is differentiated from invasive mole (chorio-adenoma destruens) by:

Options :

- ✘ presence of high titre of urinary chorionic gonadotropins
- ✘ presence of cannon ball shadow in lungs
- ✔ absence of villi structure on histological examination of the lesion
- ✘ All of the given options are correct

Question Number : 122 Question Id : 57996510224 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A positive "stallworthy's sign " is suggestive of which of the following conditions :

Options :

- ✘ twin pregnancy
- ✘ breech presentation
- ✘ vesicular mole
- ✔ low lying placenta

Question Number : 123 Question Id : 57996510225 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The following test may be abnormal in disseminated intravascular coagulation except

Options :

- ✘ prothrombin
- ✘ activated partial thromboplastin time
- ✘ D-timer levels
- ✔ clot solubility

Question Number : 124 Question Id : 57996510226 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The following test are related to blood coagulation disorder in obstetrics except

Options :

- ✔ thrombocytopenia is a feature of fibrinolytic process and not of DIC
- ✘ in DIC RBC will be 'helmet shaped or fragmented, but in fibrinolytic process the cell morphology is normal
- ✘ Weiner clot observation test gives a rough estimation of total blood fibrinogen level
- ✘ thrombocytopenia can be diagnosed from the peripheral smear

Question Number : 125 Question Id : 57996510227 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common cause of postpartum hemorrhage is:

Options :

1. ✓ uterine atony
2. ✗ Retained products
3. ✗ trauma
4. ✗ Bleeding disorders

Question Number : 126 Question Id : 57996510228 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True about placenta accreta is

Options :

1. ✓ seen in cesarean scar
2. ✗ removal should be done under GA in piecemeal
3. ✗ chorionic villi invade serosa
4. ✗ it is an etiological factor for amniotic fluid embolism

Question Number : 127 Question Id : 57996510229 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Best indicator of Thyroid function in pregnancy is

Options :

1. ✗ TBG
2. ✗ TSH
3. ✓ Free T4
4. ✗ Anti Thyroid antibody

Question Number : 128 Question Id : 57996510230 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

A patient went into shock immediately after normal delivery, likely cause is:

Options :

1. ✗ amniotic fluid embolism
2. ✗ PPH
3. ✓ uterine inversion
4. ✗ eclampsia

Question Number : 129 Question Id : 57996510231 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Common cause of retained placenta

Options :

1. ✓ atonic uterus
2. ✗ constriction ring
3. ✗ placenta accrete
4. ✗ poor voluntary expulsive effort

Question Number : 130 Question Id : 57996510232 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most common type of twin pregnancy is

Options :

1. ✘ vertex + transverse
2. ✔ both vertex
3. ✘ vertex + breech
4. ✘ both breech

Question Number : 131 Question Id : 57996510233 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

To say twin discordance the difference in the two twins should be

Options :

1. ✘ 15% with the larger twin as index
2. ✘ 15% with the smaller twin as index
3. ✔ 25% with the larger twin as index
4. ✘ 25% with the smaller twin as index

Question Number : 132 Question Id : 57996510234 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Correct statement about establishing the chorionicity in twin pregnancy is :

Options :

1. ✘ same sex rule out dichorionicity
2. ✔ twin peak in dichorionicity
3. ✘ thick membrane is present in monochorionic
4. ✘ best detected after 16 weeks

Question Number : 133 Question Id : 57996510235 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Vaginal delivery is allowed in all except

Options :

1. ✔ monochorionic monoamniotic twins
2. ✘ first twin cephalic and second breech
3. ✘ extended breech
4. ✘ mento anterior

Question Number : 134 Question Id : 57996510236 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Blood chimerism is maintained by:

Options :

1. ✔ monochorionic dizygotic twins
2. ✘ dichorionic dizygotic twins
3. ✘ vanishing twins
4. ✘ singleton pregnancy

Question Number : 135 Question Id : 57996510237 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Monochorionic monoamniotic twin occurs if division occurs

Options :

1. ✘ before 24 hours
2. ✘ 1-4 days
3. ✘ 4-8 days
4. ✔ >8 days

Question Number : 136 Question Id : 57996510238 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Drug of choice to reduce polyhydramnios is

Options :

1. ✘ Aspirin
2. ✔ Indomethacin
3. ✘ Frusemide
4. ✘ Misoprostol

Question Number : 137 Question Id : 57996510239 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Absolute proof of monozygosity is determined by

Options :

1. ✔ DNA finger printing
2. ✘ intervening membrane layers
3. ✘ sex of the babies
4. ✘ reciprocal skin grafting

Question Number : 138 Question Id : 57996510240 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which of the following test is most sensitive for the detection of iron depletion in pregnancy?

Options :

1. ✘ serum iron
2. ✔ serum ferritin
3. ✘ serum transferrin
4. ✘ serum iron binding capacity

Question Number : 139 Question Id : 57996510241 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In pregnancy, which type of anemia is not common in India

Options :

1. ✔ Vitamin B12 anemia
2. ✘ folic acid anemia
3. ✘ iron + folic acid anemia
4. ✘ iron deficiency anemia

Question Number : 140 Question Id : 57996510242 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The following statement are related to the treatment of thalassaemia except

Options :

1. ✘ fresh (relatively) blood transfusion
2. ✘ folic acid
3. ✔ routine iron therapy
4. ✘ deferoxamine improves pregnancy outcome

Question Number : 141 Question Id : 57996510243 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Formula used for estimation of the total iron requirement is

Options :

1. ✘ $4 \times \text{body weight (kg)} \times \text{Hb deficit (g/dl)}$
2. ✔ $4.4 \times \text{body weight(kg)} \times \text{Hb deficit (g/dl)}$
3. ✘ $0.3 \times \text{body weight(kg)} \times \text{Hb deficit (g/dl)}$
4. ✘ $3.3 \times \text{body weight(kg)} \times \text{Hb deficit(g/dl)}$

Question Number : 142 Question Id : 57996510244 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Maximum cardiac output in pregnancy is at

Options :

1. ✘ 20 weeks
2. ✘ 24 weeks
3. ✘ 26 weeks
4. ✔ 28 weeks

Question Number : 143 Question Id : 57996510245 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Indication for Internal Pelvic version in modern obstetrics is

Options :

1. ✘ Oblique line
2. ✘ Shoulder presentation
3. ✔ Second of Transverse lie
4. ✘ Mento posterior

Question Number : 144 Question Id : 57996510246 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

In Heart patient the worst prognosis during pregnancy is seen in

Options :

1. ✘ mitral regurgitation
2. ✘ mitral valve prolapse
3. ✔ aortic stenosis
4. ✘ pulmonary stenosis

Question Number : 145 Question Id : 57996510247 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Patient with organic heart disease in pregnancy most commonly die during

Options :

1. ✘ 20 -24 weeks of pregnancy
2. ✘ first stage of labour
3. ✔ soon following delivery
4. ✘ two weeks postpartum

Question Number : 146 Question Id : 57996510248 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

True about diabetes in pregnancy are all except

Options :

1. ✘ glucose challenge test is done between 24-28 weeks
2. ✘ 50 gm of sugar is given for screening test
3. ✔ insulin resistance improves with pregnancy
4. ✘ diabetes control before conception is important to prevent malformation

Question Number : 147 Question Id : 57996510249 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Most sensitive screening test in diabetic mother for congenital malformation is

Options :

1. ✘ MS AFP
2. ✘ blood glucose
3. ✘ amniotic fluid AFP
4. ✔ Hb A1C (glycosylated hemoglobin)

Question Number : 148 Question Id : 57996510250 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Fasting blood sugar in diabetic female in any stage of pregnancy should be maintained as

Options :

1. ✔ 70-100 mg %
2. ✘ 100-130 mg%
3. ✘ 130-160 mg%
4. ✘ 160-190 mg%

Question Number : 149 Question Id : 57996510251 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

The commonest congenital malformation in infant of a diabetic mother is

Options :

1. ✘ multicystic kidneys
2. ✘ Esophageal atresia
3. ✔ Neural tube defect
4. ✘ duodenal atresia

Question Number : 150 Question Id : 57996510252 Question Type : MCQ Option Shuffling : Yes Display Question Number : Yes
Single Line Question Option : No Option Orientation : Vertical

Correct : 1 Wrong : 0.33

Which type of eclampsia has the worst prognosis:

Options :

1. ✓ Anti partum
2. ✗ Post Partum.
3. ✗ Intra Partum.
4. ✗ Imminent.