

Question Paper Preview

Notations :

- Options shown in green color and with ✓ icon are correct.
- Options shown in red color and with ✗ icon are incorrect.

Subject Name:	Dental Surgery1
Calculator:	Scientific
Magnifying Glass Required?:	No
Ruler Required?:	No
Eraser Required?:	No
Scratch Pad Required?:	No
Rough Sketch/Notepad Required?:	No
Protractor Required?:	No

Display Number Panel:	Yes
Group All Questions:	No

Question Number : 1 Question Id : 10922233581 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The carotid canal transmits all the following except

Options :

- ✗ Internal carotid artery
- ✓ External carotid artery
- ✗ Sympathetic plexus
- ✗ Venous plexus

Question Number : 2 Question Id : 10922233582 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Bleeding and discharge of CSF through the ear occurs commonly with fracture of

Options :

- ✗ Anterior cranial fossa
- ✓ Middle cranial fossa
- ✗ Posterior cranial fossa
- ✗ None of the given options

Question Number : 3 Question Id : 10922233583 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Part of the skull that is more fragile

Options :

- ✓ Base of the skull
- ✗ Vault of the skull

- ✘ Lateral aspect of the skull
- ✘ Posterior aspect of the skull

Question Number : 4 Question Id : 10922233584 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The skeletal derivatives of Reichert's cartilage are

Options :

- ✘ Greater cornua of hyoid and lower half of hyoid bone
- ✘ Thyroid, cricoid, arytenoid, cuneiform, corniculate cartilage
- ✔ Stapes, styloid process, stylohyoid ligament, lesser cornua
- ✘ Maxilla, zygomatic, squamous part of temporal bone

Question Number : 5 Question Id : 10922233585 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Dangerous area of the scalp is formed by

Options :

- ✘ Skin
- ✘ Epicranial aponeurosis
- ✔ Loose areolar tissue
- ✘ Epicranium

Question Number : 6 Question Id : 10922233586 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The muscle that acts as a sphincter for the nostrils is

Options :

- ✘ Dilator naris
- ✘ Depressor septi
- ✘ Levator labii superioris alaeque nasi
- ✔ Compressor naris

Question Number : 7 Question Id : 10922233587 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Paralysis in only lower part of the face is caused by

Options :

- ✘ Infranuclear lesions
- ✔ Supranuclear lesions
- ✘ Bells palsy
- ✘ Infranuclear lesions or Supranuclear lesions

Question Number : 8 Question Id : 10922233588 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Secretomotor fibers reach the parotid gland through ----- nerve.

Options :

- ✘ Glossopharyngeal
- ✘ Tympanic
- ✘ Lesser petrosal
- ✔ Auriculotemporal

Question Number : 9 Question Id : 10922233589 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Branches given by facial nerve within the canal include all the following EXCEPT

Options :

1. ✘ Greater petrosal nerve
2. ✘ Chorda tympani
3. ✔ Posterior auricular
4. ✘ Nerve to stapedius

Question Number : 10 Question Id : 10922233590 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

An accessory ligament of TMJ that represents thickened part of deep cervical fascia that separates parotid and submandibular gland is

Options :

1. ✘ Capsular ligament
2. ✘ Lateral temporomandibular ligament
3. ✘ Sphenomandibular ligament
4. ✔ Stylomandibular ligament

Question Number : 11 Question Id : 10922233591 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Motor part of mandibular nerve is tested by asking the patient to

Options :

1. ✘ Clench teeth
2. ✘ Move the lower jaw from side to side
3. ✘ Open the mouth
4. ✔ All the given options

Question Number : 12 Question Id : 10922233592 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Thyroid gland is located against ----- cervical vertebrae

Options :

1. ✔ C5, C6, C7
2. ✘ C4, C5, C6
3. ✘ C2, C3, C4
4. ✘ C3, C4, C5

Question Number : 13 Question Id : 10922233593 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The superior parathyroid glands develop from

Options :

1. ✘ Third pharyngeal pouch
2. ✔ Fourth pharyngeal pouch
3. ✘ Second pharyngeal pouch
4. ✘ Sixth pharyngeal pouch

Question Number : 14 Question Id : 10922233594 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Tonsil and tongue drain into ----- lymph nodes respectively

Options :

1. ✘ Jugulo-omohyoid & anterior cervical

2. ✓ Jugulo-digatric & jugulo-omohyoid
3. ✗ Waldeyers ring and jugulo-omohyoid
4. ✗ Prelaryngeal & jugulo-digatric

Question Number : 15 Question Id : 10922233595 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Tick the correct statement

Options :

1. ✗ All the muscles of soft palate are supplied by pharyngeal plexus
2. ✗ All the muscles except levator veli palatini are supplied by mandibular nerve
3. ✓ All the muscles except tensor veli palatini are supplied by pharyngeal plexus
4. ✗ Only palatoglossus & palatopharyngeus are supplied by vagus nerve

Question Number : 16 Question Id : 10922233596 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Maxillary sinus opens into

Options :

1. ✗ Anterior part of hiatus semilunaris
2. ✓ Posterior part of hiatus semilunaris
3. ✗ Inferior meatus
4. ✗ Sphenoethmoidal recess

Question Number : 17 Question Id : 10922233597 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Paralysis of tongue muscles associated with atrophy of one half of the tongue is seen in

Options :

1. ✓ Infra nuclear lesion / injury to hypoglossal nerve
2. ✗ Supranuclear lesion / injury to hypoglossal nerve
3. ✗ Pseudobulbar paralysis
4. ✗ All the given options

Question Number : 18 Question Id : 10922233598 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Identify the correct relation.

Options :

1. ✗ Jaw jerk reflex : Mandibular division
2. ✗ Sneeze reflex: Maxillary division
3. ✗ Corneal blink reflex: Ophthalmic division
4. ✓ All the given options

Question Number : 19 Question Id : 10922233599 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Pterygoid plexus communicates with

Options :

1. ✗ Cavernous sinus through emissary veins
2. ✗ Inferior ophthalmic vein through inferior orbital fissure
3. ✗ Facial vein through deep facial vein
4. ✓ All the given options

Question Number : 20 Question Id : 10922233600 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Pulsating exophthalmos is caused by communication between

Options :

1. ✘ sigmoid sinus and mastoid vein
2. ✘ transverse sinus & posterior temporal vein
3. ✔ cavernous sinus & internal carotid artery
4. ✘ straight sinus & dangerous area of the face

Question Number : 21 Question Id : 10922233601 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Torniquet test is positive in

Options :

1. ✔ Vitamin C deficiency
2. ✘ Vitamin B 12 deficiency
3. ✘ Iron deficiency
4. ✘ Zinc deficiency

Question Number : 22 Question Id : 10922233602 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

After birth, RBC are almost exclusively formed in

Options :

1. ✔ Bone marrow
2. ✘ Spleen
3. ✘ Liver
4. ✘ Yolk sac

Question Number : 23 Question Id : 10922233603 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Effects of anemia includes

Options :

1. ✘ Increases resistance to blood flow in tissues
2. ✘ Increases viscosity of blood
3. ✔ Greater pumping load on heart
4. ✘ Decreases cardiac output

Question Number : 24 Question Id : 10922233604 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Function of basophils is

Options :

1. ✘ Phagocytosis
2. ✘ Antibody formation
3. ✘ Formation of reactive oxygen species
4. ✔ Release of histamine, bradykinin, serotonin

Question Number : 25 Question Id : 10922233605 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Low serum calcium concentration promotes the secretion of

Options :

1. ✘ Thyroxine

2. ✘ ACTH
3. ✔ PTH
4. ✘ Insulin

Question Number : 26 Question Id : 10922233606 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Metabolic effects of growth hormone include

Options :

1. ✘ Increased rate of protein synthesis
2. ✘ Increased mobilization of fats from adipose tissue
3. ✔ Decreased rate of peripheral utilisation of glucose
4. ✘ All the given options

Question Number : 27 Question Id : 10922233607 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Iodide in high concentration (100 times normal plasma concentration) leads to

Options :

1. ✘ Increases endocytosis of colloid from follicles
2. ✘ Increases thyroid gland size
3. ✘ Causes hyperthyroidism
4. ✔ Decreases thyroid activity

Question Number : 28 Question Id : 10922233608 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Type 1 and type 2 diabetes may be distinguished by measuring

Options :

1. ✘ Blood glucose
2. ✘ Urinary glucose
3. ✘ Plasma Insulin
4. ✔ All the given options

Question Number : 29 Question Id : 10922233609 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Complications of diabetes include all the following except

Options :

1. ✘ Atherosclerosis
2. ✘ Increased susceptibility to infection
3. ✘ Cataract formation
4. ✔ Obesity

Question Number : 30 Question Id : 10922233610 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Vascular haemophilia occurs due to the deficiency of

Options :

1. ✘ Factor VIII
2. ✘ Factor IX
3. ✔ von Willebrand factor
4. ✘ Fibrinogen

Question Number : 31 Question Id : 10922233611 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Volume of air normally inspired or expired with each breath is

Options :

- ✘ Vital capacity
- ✔ Tidal volume
- ✘ Functional reserve volume
- ✘ Residual volume

Question Number : 32 Question Id : 10922233612 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Pulmonary edema may be caused by

Options :

- ✘ Left sided heart failure
- ✘ Infections
- ✘ Breathing noxious gases
- ✔ All the given options

Question Number : 33 Question Id : 10922233613 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Asthmatic patients have difficulty during

Options :

- ✘ Inspiration
- ✔ Expiration
- ✘ Inspiration and Expiration
- ✘ None of the given options

Question Number : 34 Question Id : 10922233614 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Hypercapnia is caused by

Options :

- ✘ All forms of hypoxia
- ✔ Hypoxia caused by hypoventilation
- ✘ Hypoxia caused due to anemia
- ✘ Hypoxia due to poisoning of oxidative enzymes

Question Number : 35 Question Id : 10922233615 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

During pharyngeal phase, respiration

Options :

- ✔ Is halted
- ✘ Continues to occur
- ✘ May or may not occur
- ✘ Varies with the bolus size

Question Number : 36 Question Id : 10922233616 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Hormones increasing intestinal motility include all the following except

Options :

- ✔ Glucagon

2. ✘ Gastrin
3. ✘ Serotonin
4. ✘ CCK

Question Number : 37 Question Id : 10922233617 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Salivary stimulation is caused by

Options :

1. ✘ Parasympathetic nervous stimulation
2. ✘ Reflexes of stomach & upper intestines
3. ✘ Sympathetic stimulation
4. ✔ All the given options

Question Number : 38 Question Id : 10922233618 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Phase/phases of gastric secretion include

Options :

1. ✘ Cephalic phase
2. ✘ Gastric phase
3. ✘ Intestinal phase
4. ✔ All the given options

Question Number : 39 Question Id : 10922233619 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Causes for peptic ulcer disease include all the following except

Options :

1. ✘ Consumption of alcohol
2. ✘ Tobacco smoking
3. ✘ Infection by Helicobacter pylori
4. ✔ Hypochlorhydria

Question Number : 40 Question Id : 10922233620 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Inability to perceive a particular taste is called

Options :

1. ✘ Taste resistance
2. ✘ Taste aversion
3. ✔ Taste blindness
4. ✘ Cacogeusia

Question Number : 41 Question Id : 10922233621 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Highest modulus of elasticity is seen with

Options :

1. ✔ Enamel
2. ✘ Dentin
3. ✘ Composite
4. ✘ Amalgam

Question Number : 42 Question Id : 10922233622 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The restorative material with coefficient of thermal expansion closest to enamel is

Options :

- GIC
- Silicate cement
- Composite
- Polycarboxylate cement

Question Number : 43 Question Id : 10922233623 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Impression material used to make impressions of partial edentulous mouth are

Options :

- Impression compound
- Impression wax
- Hydrocolloid
- None of the given options

Question Number : 44 Question Id : 10922233624 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Rubber base impressions are poured immediately as they

Options :

- Become elastic
- Continue to polymerize
- Show gelation
- Show imbibition

Question Number : 45 Question Id : 10922233625 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Corrective impression is made by

Options :

- Impression paste
- Impression plaster
- Alginate
- Rubber base impression materials

Question Number : 46 Question Id : 10922233626 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Crosslinking agent of polysulphide impression material is

Options :

- Stannous octate
- Sulfonate ester
- Lead dioxide
- Platinum

Question Number : 47 Question Id : 10922233627 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Impression material that do not set/harden by chemical reaction is

Options :

- Alginate

2. ✘ ZOE paste
3. ✔ Impression compound
4. ✘ Plaster of paris

Question Number : 48 Question Id : 10922233628 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Setting time in Plaster of Paris is most effectively altered by

Options :

1. ✘ Altering temperature of water
2. ✔ Addition of accelerators and retarders
3. ✘ Speed and length of hand spatulation
4. ✘ Altering powder /liquid ratio

Question Number : 49 Question Id : 10922233629 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Internal porosity in thick areas of denture is caused by

Options :

1. ✘ Insufficient pressure or
2. ✘ Voids due to air inclusion
3. ✘ Inadequate material
4. ✔ Curing temperature more than 100 degree C

Question Number : 50 Question Id : 10922233630 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Etching agent is

Options :

1. ✘ Methyl methacrylate
2. ✔ Phosphoric acid
3. ✘ UV light
4. ✘ Eugenol

Question Number : 51 Question Id : 10922233631 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Which factor improves in acrylic denture bases containing cross linking agents

Options :

1. ✘ Surface hardness
2. ✔ Craze resistance
3. ✘ Tissue compatibility
4. ✘ Colour

Question Number : 52 Question Id : 10922233632 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Packing of acrylic resin mixture should be done in

Options :

1. ✔ Doughy stage
2. ✘ Sandy stage
3. ✘ Stringy stage
4. ✘ Sticky stage

Question Number : 53 Question Id : 10922233633 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Residual monomer content in heat cured acrylic is

Options :

1. ✘ 3- 5%
2. ✘ 0.3-0.5%
3. ✘ 2-5%
4. ✔ 0.2-0.5%

Question Number : 54 Question Id : 10922233634 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Composition of amalgam alloy includes

Options :

1. ✘ 65% silver, 32% tin, 3% copper
2. ✘ 49% silver, 32%tin, 19%copper
3. ✔ 65%silver, 29% tin, 5-6%copper
4. ✘ 50% silver, 25% tin, 25% copper

Question Number : 55 Question Id : 10922233635 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

High copper amalgam is superior to other amalgam due to

Options :

1. ✔ Less marginal breakdown
2. ✘ Higher ratio of tensile to compressive strength
3. ✘ Workability at lower Hg-alloy ratio
4. ✘ Have less resistance to tarnish & corrosion

Question Number : 56 Question Id : 10922233636 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

NiTi alloys have

Options :

1. ✘ Corrosion resistance
2. ✔ Shape memory
3. ✘ Hyperrigidity
4. ✘ All the given options

Question Number : 57 Question Id : 10922233637 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Stainless steel orthodontic wire can be hardened by

Options :

1. ✘ Tempering
2. ✔ Work hardening
3. ✘ Age hardening
4. ✘ Precipitation hardening

Question Number : 58 Question Id : 10922233638 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Incomplete casting can result from

Options :

1. ✘ Narrow sprue diameter

2. ✘ High viscosity of the metal
3. ✘ Incomplete wax pattern elimination
4. ✔ All the given options

Question Number : 59 Question Id : 10922233639 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Most rigid material is

Options :

1. ✘ Co-Ni-Cr
2. ✘ Wrought Ni-Cr
3. ✘ Wrought Cr-Co
4. ✔ Cast Cr-Co

Question Number : 60 Question Id : 10922233640 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Pickling solution used for gypsum bonded investments is

Options :

1. ✔ HCl
2. ✘ HNO₃
3. ✘ H₂SO₄
4. ✘ H₃PO₄

Question Number : 61 Question Id : 10922233641 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Best compatibility with pulp is seen with

Options :

1. ✘ ZOE
2. ✔ GIC
3. ✘ Silicate cement
4. ✘ Zinc phosphate cement

Question Number : 62 Question Id : 10922233642 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

In a PFM, porcelain binds to metal by

Options :

1. ✘ Metallic bond
2. ✘ Mechanical bond
3. ✘ Chemical bond
4. ✔ Mechanical bond and Chemical bond

Question Number : 63 Question Id : 10922233643 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Main constituent of porcelain

Options :

1. ✘ Kaolin
2. ✘ Clay
3. ✔ Feldspar
4. ✘ Quartz

Question Number : 64 Question Id : 10922233644 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Outer and inner epithelium first appear in

Options :

1. ✘ Bud stage
2. ✔ Cap stage
3. ✘ Bell stage
4. ✘ Advanced bell stage

Question Number : 65 Question Id : 10922233645 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Enamel does not have self repair capacity as

Options :

1. ✘ It is mostly inorganic
2. ✔ Formative cells are lost once enamel is completely formed
3. ✘ Has no blood supply
4. ✘ Stretched far away from pulp

Question Number : 66 Question Id : 10922233646 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Main bulk of dentin is formed by

Options :

1. ✘ Mantle dentin
2. ✘ Predentin
3. ✔ Intertubular dentin
4. ✘ Peritubular dentin

Question Number : 67 Question Id : 10922233647 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Aspiration of odontoblasts into dentinal tubules occurs due to

Options :

1. ✔ Cavity preparation
2. ✘ Dehydration
3. ✘ Irrigation
4. ✘ None of the given options

Question Number : 68 Question Id : 10922233648 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Interdependency of osteoblasts and osteoclasts in bone remodelling is known as

Options :

1. ✘ Interaction
2. ✘ Interdependency
3. ✔ Coupling
4. ✘ Symbiosis

Question Number : 69 Question Id : 10922233649 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Resistance to masticatory forces is given by ----- fibres of periodontal ligament

Options :

1. ✘ Alveolar crest

2. ✘ Transeptal
3. ✘ Apical
4. ✔ Oblique

Question Number : 70 Question Id : 10922233650 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Calcification of lamina dura when compared to alveolar bone is -----

Options :

1. ✔ Same
2. ✘ Lesser
3. ✘ Greater
4. ✘ Varies from place to place

Question Number : 71 Question Id : 10922233651 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

TMJ starts developing at an age of

Options :

1. ✘ 6 weeks
2. ✘ 8 weeks
3. ✔ 10 weeks
4. ✘ 20 weeks

Question Number : 72 Question Id : 10922233652 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Type A cells of synovial membrane of TMJ are

Options :

1. ✘ Fibroblast like
2. ✔ Macrophage like
3. ✘ Lymphocyte like
4. ✘ Inbetween cell type A & B

Question Number : 73 Question Id : 10922233653 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Mushroom shaped papillae that contain few taste buds are

Options :

1. ✘ Circumvallate papillae
2. ✔ Fungiform papillae
3. ✘ Filiform papillae
4. ✘ Foliate papillae

Question Number : 74 Question Id : 10922233654 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The non keratinocytes of oral epithelium include all the following except

Options :

1. ✘ Langerhans cells
2. ✘ Melanocytes
3. ✔ Spinous cells
4. ✘ Merkel cells

Question Number : 75 Question Id : 10922233655 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Gingiva is predominantly

Options :

1. ✘ Orthokeratinised
2. ✔ Parakeratinised
3. ✘ Nonkeratinised
4. ✘ All the given options

Question Number : 76 Question Id : 10922233656 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Eruption of permanent teeth is facilitated by

Options :

1. ✘ Exfoliation of primary teeth
2. ✔ Formation of gubernacular canal
3. ✘ Osteoclastic activity
4. ✘ All the given options

Question Number : 77 Question Id : 10922233657 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Ectopic sebaceous glands in the oral cavity are known as

Options :

1. ✘ Koplik spots
2. ✔ Fordyce's spots
3. ✘ Rose spots
4. ✘ Bitot spots

Question Number : 78 Question Id : 10922233658 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Herpetic ulcers and aphthous ulcers have predilection for

Options :

1. ✘ Non keratinised mucosa
2. ✘ Keratinised mucosa
3. ✔ Keratinised mucosa & Non keratinised mucosa
4. ✘ Non keratinised mucosa & Keratinised mucosa

Question Number : 79 Question Id : 10922233659 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Supernumerary teeth are formed due to disturbance at

Options :

1. ✔ Initiation stage
2. ✘ Apposition stage
3. ✘ Histodifferentiation stage
4. ✘ Morphodifferentiation stage

Question Number : 80 Question Id : 10922233660 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

First formed dentin that is mineralised in a tooth is called as

Options :

1. ✘ Predentin

2. ✓ Mantle dentin
3. ✗ Secondary dentin
4. ✗ Circumpulpal dentin

Question Number : 81 Question Id : 10922233661 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Toxigenicity and drug resistance of bacteria is conferred by

Options :

1. ✗ Voluntin granules
2. ✓ Plasmids
3. ✗ Capsule
4. ✗ Mesosomes

Question Number : 82 Question Id : 10922233662 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

A person who had recovered from the disease but continues to harbour the infection are known as

Options :

1. ✗ Contact carrier
2. ✗ Paradoxical carrier
3. ✓ Convalescent carrier
4. ✗ Subclinical carrier

Question Number : 83 Question Id : 10922233663 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

About 80% of serum immunoglobulins are constituted by

Options :

1. ✗ IgA
2. ✗ IgM
3. ✓ IgG
4. ✗ IgE

Question Number : 84 Question Id : 10922233664 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The medium used to differentiate lactose and non lactose fermenters is

Options :

1. ✗ Nutrient agar
2. ✗ Blood agar
3. ✓ Mac Conkey agar
4. ✗ Thioglycolate medium

Question Number : 85 Question Id : 10922233665 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Systemic lupus erythematosus is due to ----- hypersensitivity reaction

Options :

1. ✗ Type I
2. ✗ Type II
3. ✓ Type III
4. ✗ Type IV

Question Number : 86 Question Id : 10922233666 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Bactericidal drugs are most effective at which phase of bacterial growth?

Options :

1. ✘ Lag phase
2. ✔ Log phase
3. ✘ Zero growth phase
4. ✘ Phase of decline

Question Number : 87 Question Id : 10922233667 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Which of the following antigens represent active intrahepatic viral replication of HBV

Options :

1. ✘ HBsAg
2. ✘ HBcAg
3. ✔ HBeAg
4. ✘ All the given options

Question Number : 88 Question Id : 10922233668 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Oncogenic viruses include all the following except

Options :

1. ✘ HSV
2. ✘ HPV
3. ✘ EBV
4. ✔ VZV

Question Number : 89 Question Id : 10922233669 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Test to identify HIV infection during window period is

Options :

1. ✘ Western Blot
2. ✘ ELISA
3. ✔ p24 antigen
4. ✘ VDRL

Question Number : 90 Question Id : 10922233670 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Candida is

Options :

1. ✘ Opportunistic fungal infection
2. ✘ Co-carcinogen
3. ✔ Opportunistic fungal infection and Co-carcinogen
4. ✘ None of the given options

Question Number : 91 Question Id : 10922233671 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

When a graft is taken from an identical twin, it is known as

Options :

1. ✘ Autograft

2. ✓ Isograft
3. ✗ Allograft
4. ✗ Xenograft

Question Number : 92 Question Id : 10922233672 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Gas gangrene is caused by

Options :

1. ✗ Clostridium tetani
2. ✓ Clostridium perfringens
3. ✗ Clostridium difficile
4. ✗ Clostridium botulinum

Question Number : 93 Question Id : 10922233673 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The enzymes secreted by Streptococcus species responsible for spread of infection are

Options :

1. ✗ Pyogenic toxin
2. ✓ Hyaluronidase
3. ✗ Streptodornase
4. ✗ Deoxyribonuclease

Question Number : 94 Question Id : 10922233674 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Step ladder pyrexia and rose spots occur in

Options :

1. ✓ Enteric fever
2. ✗ Tuberculosis
3. ✗ Tetanus
4. ✗ Syphilis

Question Number : 95 Question Id : 10922233675 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Positive tuberculin test indicates

Options :

1. ✗ Infection with tubercle bacillus
2. ✗ BCG immunization
3. ✓ Infection with tubercle bacillus and BCG immunization
4. ✗ None of the given options

Question Number : 96 Question Id : 10922233676 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

100% bioavailability of drugs is achieved by ----- route of administration

Options :

1. ✗ Oral
2. ✗ Inhalation
3. ✓ Intravenous
4. ✗ Transcutaneous

Question Number : 97 Question Id : 10922233677 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Systemic effects produced by rubbing a drug onto the skin is called

Options :

1. ✘ Iontophoresis
2. ✔ Inunction
3. ✘ Jet injection
4. ✘ Intrathecal administration

Question Number : 98 Question Id : 10922233678 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

First order kinetics of drug elimination includes

Options :

1. ✔ Constant fraction of drug disappears in equal intervals of time at a rate proportional to its plasma concentration
2. ✘ Fixed quantity of drug is eliminated per unit time
3. ✘ Only half the drug is eliminated, rest is retained
4. ✘ Elimination is increased by drug metabolism in liver

Question Number : 99 Question Id : 10922233679 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Therapeutic index is

Options :

1. ✘ Median effective dose/ Median lethal dose
2. ✔ Median lethal dose/ Median effective dose
3. ✘ Median effective dose/ Mean number of side effects in test population
4. ✘ Median lethal dose/ Mean number of side effects in test population

Question Number : 100 Question Id : 10922233680 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Antagonism caused by 2 agonists acting independently but causing opposite actions is called as

Options :

1. ✘ Competitive antagonism
2. ✘ Chemical antagonism
3. ✔ Functional antagonism
4. ✘ Non-competitive antagonism

Question Number : 101 Question Id : 10922233681 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

All the following antibiotics act by inhibiting protein synthesis except

Options :

1. ✔ Vancomycin
2. ✘ Tetracycline
3. ✘ Erythromycin
4. ✘ Chloramphenicol

Question Number : 102 Question Id : 10922233682 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Antibiotics concentrated in gingival crevice and preferred for periodontal diseases are

Options :

1. ✘ Penicillins
2. ✔ Tetracyclines
3. ✘ Cephalosporins
4. ✘ Macrolides

Question Number : 103 Question Id : 10922233683 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Gray baby syndrome in infants is caused by administration of

Options :

1. ✔ Chloramphenicol in infants lacking glucuronyl transferase
2. ✘ Tetracycline in infants lacking alanine transferase
3. ✘ Penicillin in children lacking streptokinase
4. ✘ Macrolides in children lacking cholinesterase

Question Number : 104 Question Id : 10922233684 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Tuberculocidal drugs safe in pregnancy are

Options :

1. ✘ Streptomycin, INH
2. ✘ Rifampicin, Ethambutal
3. ✘ INH, Pyrazinamide
4. ✔ INH, Ethambutal

Question Number : 105 Question Id : 10922233685 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Pseudomembranous colitis caused by Clindamycin is treated by

Options :

1. ✘ Actinomycin D
2. ✘ Erythromycin
3. ✔ Vancomycin
4. ✘ Lincomycin

Question Number : 106 Question Id : 10922233686 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Neurotransmitter released at postganglionic sympathetic nerves is

Options :

1. ✘ Acetylcholine
2. ✘ Adrenaline
3. ✔ Noradrenaline
4. ✘ GABA

Question Number : 107 Question Id : 10922233687 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Adrenaline is the drug of choice in anaphylactic shock as

Options :

1. ✔ It increases BP and relieves bronchospasm
2. ✘ It decreases BP and causes bronchodilation
3. ✘ It increases BP and causes bronchoconstriction
4. ✘ It decreased BP and causes bronchoconstriction

Question Number : 108 Question Id : 10922233688 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Morphine is indicated for all the following conditions except

Options :

1. ✘ Severe visceral pain
2. ✘ Pain due to cancer
3. ✔ Head injury
4. ✘ Preanesthetic medication

Question Number : 109 Question Id : 10922233689 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Local anesthetic consists of

Options :

1. ✘ Aromatic lipophilic group
2. ✘ Hydrophilic amino group
3. ✘ Intermediate chain(Ester / Amide)
4. ✔ All the given options

Question Number : 110 Question Id : 10922233690 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Tannic acid given in gingival diseases acts as

Options :

1. ✔ Astringent
2. ✘ Disinfectant
3. ✘ Sterilizing agent
4. ✘ All the given options

Question Number : 111 Question Id : 10922233691 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Clavulanic acid is added to amoxicillin to

Options :

1. ✘ Increase the half life of amoxicillin
2. ✔ Increase antibacterial spectrum of amoxicillin
3. ✘ Increase the bioavailability of amoxicillin
4. ✘ Reduce the toxicity of amoxicillin

Question Number : 112 Question Id : 10922233692 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Mucocutaneous candidiasis may be treated by

Options :

1. ✘ Amphotericin B
2. ✘ Ketoconazole
3. ✘ Fluconazole
4. ✔ All the given options

Question Number : 113 Question Id : 10922233693 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Antiretroviral treatment regime suggested for post exposure prophylaxis for HIV is

Options :

1. ✓ Zidovudine, Lamivudine, Lopinavir
2. ✗ Amoxicillin /ampicillin / Erythromycin
3. ✗ HBIG
4. ✗ All the given options

Question Number : 114 Question Id : 10922233694 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Patient on antitubercular therapy for around 3 months developed tingling of extremities. He should be treated with

Options :

1. ✗ Gabapentine
2. ✓ Pyridoxine
3. ✗ Cyanocobalamine
4. ✗ Cannot be treated

Question Number : 115 Question Id : 10922233695 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Drug of choice in osteomyelitis is

Options :

1. ✗ Cephalexin
2. ✗ Penicillin
3. ✓ Clindamycin
4. ✗ Ciprofloxacin

Question Number : 116 Question Id : 10922233696 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

When the mesiobuccal cusp of maxillary first molar falls into the interdental embrasure between mandibular first & second molar it is

Options :

1. ✗ Class I molar relation
2. ✗ Class II molar relation
3. ✓ Class III molar relation
4. ✗ Class IV molar relation

Question Number : 117 Question Id : 10922233697 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Process of destruction of all microbial forms is

Options :

1. ✗ Asepsis
2. ✓ Sterilization
3. ✗ Disinfection
4. ✗ Cleaning

Question Number : 118 Question Id : 10922233698 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Enamel rods at cervical region in primary teeth

Options :

1. ✓ Slope occlusally
2. ✗ Slope cervically

- ✘ Do not slope
- ✘ Are absent

Question Number : 119 Question Id : 10922233699 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The tooth which frequently causes palatal abscess is

Options :

- ✘ Maxillary central incisor
- ✔ Maxillary lateral incisor
- ✘ Maxillary canine
- ✘ Maxillary premolar

Question Number : 120 Question Id : 10922233700 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Albinism results from defective metabolism of

Options :

- ✔ Tyrosine
- ✘ Tryptophan
- ✘ Threonine
- ✘ Arginine

Question Number : 121 Question Id : 10922233701 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Fat soluble vitamins are

Options :

- ✔ A, D, E, K
- ✘ A,C,B, D
- ✘ A,B,C,K
- ✘ D,B,E,K

Question Number : 122 Question Id : 10922233702 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Histidine load test and FIGLU excretion test are used to estimate

Options :

- ✘ Iron
- ✔ Vitamin Folic acid
- ✘ Magnesium
- ✘ Vitamin C

Question Number : 123 Question Id : 10922233703 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Which of the following isoenzyme is elevated as the earliest sign of myocardial infarction?

Options :

- ✘ Creatine phosphokinase BB
- ✔ Creatine phosphokinase MB
- ✘ Creatine phosphokinase MM
- ✘ None of the given options

Question Number : 124 Question Id : 10922233704 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

EBV is implicated as a causative agent in all the following lesions except

Options :

1. ✘ Burkitt's lymphoma
2. ✘ Nasopharyngeal carcinoma
3. ✔ Hand, foot, mouth disease
4. ✘ Non Hodgkin's lymphoma

Question Number : 125 Question Id : 10922233705 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Verruca vulgaris spreads by

Options :

1. ✘ Contact
2. ✔ Autoinoculation
3. ✘ Droplet infection
4. ✘ Any of the above

Question Number : 126 Question Id : 10922233706 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Delayed eruption of teeth may be seen in all the following conditions except

Options :

1. ✘ Rickets
2. ✔ Mental retardation
3. ✘ Cretinism
4. ✘ Fibromatosis gingivae

Question Number : 127 Question Id : 10922233707 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

HPV positive oropharyngeal tumors have

Options :

1. ✔ Improved prognosis
2. ✘ Intermediate prognosis
3. ✘ Poor prognosis
4. ✘ Do not exist

Question Number : 128 Question Id : 10922233708 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Which tumor suppressor gene is known as guardman of the genome?

Options :

1. ✘ Rb
2. ✔ p53
3. ✘ p16
4. ✘ bcl-2

Question Number : 129 Question Id : 10922233709 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

A female in second decade presented with slight expansion of cortical plates with a multilocular radiolucency crossing the midline, anterior to premolars. Biopsy revealed giant cells with foci of hemosiderin pigment. Most likely diagnosis is

Options :

1. ✘ Tuberculous osteomyelitis

2. ✓ Central giant cell granuloma

3. ✗ Aneurysmal bone cyst

4. ✗ Osteosarcoma

Question Number : 130 Question Id : 10922233710 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Histologically, herring bone pattern is seen in

Options :

1. ✗ Cherubism

2. ✗ Multiple myeloma

3. ✓ Fibrosarcoma

4. ✗ Burkitt's lymphoma

Question Number : 131 Question Id : 10922233711 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Fluctuant, often translucent swelling of the alveolar ridge over the site of an erupting tooth is most likely

Options :

1. ✗ Dentigerous cyst

2. ✗ Eruption hematoma

3. ✗ Gingival cyst of newborn

4. ✓ Eruption cyst

Question Number : 132 Question Id : 10922233712 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Risus sardonius, opisthotonus and trismus are commonly associated with

Options :

1. ✓ Tetanus

2. ✗ Systemic sclerosis

3. ✗ Tularemia

4. ✗ Submasseteric space infection

Question Number : 133 Question Id : 10922233713 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Reactivation of herpes infection is explained by

Options :

1. ✗ Ganglionic trigger theory

2. ✗ Skin trigger theory

3. ✗ Emotional theory

4. ✓ All the given options

Question Number : 134 Question Id : 10922233714 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Fungal disease where the pathogen exhibits predilection for blood vessels and causes thrombosis of blood vessels & necrosis of tissues is

Options :

1. ✗ Candidiasis

2. ✗ Cryptococcosis

3. ✗ Rhinosporidiosis

4. ✓ Mucormycosis

Question Number : 135 Question Id : 10922233715 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Transport of metabolites and salivary buffers into the plaque and diffusion of acid out of the plaque may be affected by

Options :

1. ✘ Fructan
2. ✔ Glucan
3. ✘ Levan
4. ✘ All the given options

Question Number : 136 Question Id : 10922233716 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Heterofermentative bacteria may convert glucose into ____ acids other than lactic acid in dental plaque.

Options :

1. ✘ Propionic
2. ✘ Butyric
3. ✘ Succinic
4. ✔ All the given options

Question Number : 137 Question Id : 10922233717 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

A 11 year old patient presented clinically with a painless hard bony swelling on the outer surface of the jaw opposing a carious molar and radiologically with a onion peel appearance. The lesion most likely is

Options :

1. ✔ Proliferative periostitis
2. ✘ Exostoses
3. ✘ Enostoses
4. ✘ Fibroosseous lesion

Question Number : 138 Question Id : 10922233718 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Swelling extraorally in the region of sigmoid notch and intraorally in the tuberosity region is seen in which space infection

Options :

1. ✘ Parotid space
2. ✘ Lateral Pharyngeal space
3. ✔ Infratemporal space
4. ✘ Pterygomandibular space

Question Number : 139 Question Id : 10922233719 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Circumscribed area of tissue which is infected with exogenous pathogenic organisms, located near a mucous/cutaneous surface is called

Options :

1. ✘ Plaque
2. ✘ Focal infection
3. ✔ Focus of infection
4. ✘ Embolus

Question Number : 140 Question Id : 10922233720 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Sharp fleeting pain felt when releasing biting pressure on an object may be due to

Options :

- ✘ Trigeminal neuralgia
- ✘ Apical Periodontitis
- ✘ Pulpitis
- ✔ Cracked tooth syndrome

Question Number : 141 Question Id : 10922233721 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Mucous retention phenomenon may be seen in

Options :

- ✘ Minor & Major salivary glands
- ✘ Mucous glands of maxillary sinus
- ✔ Minor & Major salivary glands and Mucous glands of maxillary sinus
- ✘ None of the given options

Question Number : 142 Question Id : 10922233722 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Complication of healing of an extraction wound due to the placement of antibiotic ointment is called

Options :

- ✘ Dry socket
- ✔ Myospherulosis
- ✘ Osteomyelitis
- ✘ Alveolar osteitis

Question Number : 143 Question Id : 10922233723 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Stem cells of dental tissues had been identified in all the following except

Options :

- ✔ Cells of dental lamina
- ✘ Pulp of permanent teeth
- ✘ Exfoliated deciduous teeth
- ✘ Periosteum of maxillary tuberosity

Question Number : 144 Question Id : 10922233724 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Bronze diabetes occurs due to

Options :

- ✘ Insufficiency of corticosteroids
- ✔ Overload of Iron
- ✘ Oversecretion of catecholamines
- ✘ Overload of copper

Question Number : 145 Question Id : 10922233725 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Recurrent herpetic ulcers are classified under

Options :

- ✘ Recurrent Herpes Labialis

2. ✘ Recurrent Herpes Zoster
3. ✔ Recurrent Aphthous Stomatitis
4. ✘ Recurrent Erythema Multiforme

Question Number : 146 Question Id : 10922233726 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Facial paralysis is seen in all the following conditions except

Options :

1. ✘ Heerfordt's syndrome
2. ✘ Melkersson Rosenthal syndrome
3. ✘ Ramsay Hunt syndrome
4. ✔ Ascher's syndrome

Question Number : 147 Question Id : 10922233727 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Premature loss of deciduous teeth is an early manifestation of which bone disorder?

Options :

1. ✘ Osteogenesis imperfecta
2. ✔ Hypophosphatasia
3. ✘ Osteopetrosis
4. ✘ Infantile cortical hyperostosis

Question Number : 148 Question Id : 10922233728 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Bony ankyloses in which the a bony bridge extends between the ramus to zygomatic arch with an atrophic, displaced condylar head on the medial aspect is classified as

Options :

1. ✘ Type I
2. ✘ Type II
3. ✔ Type III
4. ✘ Type IV

Question Number : 149 Question Id : 10922233729 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Transmission of Hemophilia is

Options :

1. ✘ Autosomal dominant
2. ✘ Autosomal recessive
3. ✘ X linked dominant
4. ✔ X linked recessive

Question Number : 150 Question Id : 10922233730 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Tzanck cells are seen in

Options :

1. ✔ Pemphigus
2. ✘ Mucous membrane pemphigoid
3. ✘ Bullous pemphigoid
4. ✘ Lichen planus