

Question Paper Preview

Notations :

- Options shown in green color and with ✓ icon are correct.
- Options shown in red color and with ✗ icon are incorrect.

Subject Name:	Dental Surgery2
Calculator:	Scientific
Magnifying Glass Required?:	No
Ruler Required?:	No
Eraser Required?:	No
Scratch Pad Required?:	No
Rough Sketch/Notepad Required?:	No
Protractor Required?:	No

Display Number Panel:	Yes
Group All Questions:	No

Question Number : 1 Question Id : 10922234332 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Hairy tongue is due to impaired desquamation of

Options :

- ✗ Fungiform papillae
- ✓ Filiform papillae
- ✗ Circumvallate papillae
- ✗ Foliate papillae

Question Number : 2 Question Id : 10922234333 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Pemphigus vulgaris is characterised histologically by

Options :

- ✗ Sub basilar acantholysis
- ✗ Basement membrane split
- ✓ Suprabasilar acantholysis
- ✗ Subepithelial T lymphocyte band like infiltration

Question Number : 3 Question Id : 10922234334 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Differential effect of bone & soft tissue appearance on a radiograph is due to

Options :

- ✓ Higher atomic number of bone
- ✗ Higher atomic number of soft tissue

- ✘ Lesser photoelectric interaction in bone
- ✘ Greater photoelectric interactions in soft tissue

Question Number : 4 Question Id : 10922234335 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Floating teeth appearance is seen in

Options :

- ✘ Osteosarcoma
- ✘ Multiple myeloma
- ✘ Ameloblastoma
- ✔ Eosinophilic granuloma

Question Number : 5 Question Id : 10922234336 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Patient presents with a periapical radiolucency in a tooth on an IOPA taken during followup visit, treated successfully by endodontic treatment and periapical surgery. The likely reason

Options :

- ✘ Residual infection
- ✘ Periapical granuloma
- ✔ Periapical scar
- ✘ Non odontogenic cause

Question Number : 6 Question Id : 10922234337 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Radiotherapy is delivered in multiple fractions to facilitate

Options :

- ✘ Reoxygenation of the tumor
- ✘ Repair of sublethal damage in normal tissues
- ✘ Repopulation of normal tissues
- ✔ All of the given options

Question Number : 7 Question Id : 10922234338 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Ameloblastoma with best prognosis ?

Options :

- ✔ Unicystic ameloblastoma
- ✘ Solid ameloblastoma
- ✘ Granular ameloblastoma
- ✘ Mural ameloblastoma

Question Number : 8 Question Id : 10922234339 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Confirmation of a diagnosis of herpes zoster sine herpette is through

Options :

- ✘ Clinical features only
- ✘ Biopsy of the lesion
- ✔ Serology
- ✘ Imaging

Question Number : 9 Question Id : 10922234340 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Treatment of postherpetic neuralgia is

Options :

1. ✘ Capsaicin cream
2. ✘ Tricyclic antidepressants
3. ✘ Gabapentine
4. ✔ All of the given options

Question Number : 10 Question Id : 10922234341 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Abberant and uncoordinated cellular proliferation depicted at tissue level as a histologic connotation to premalignancy is termed as

Options :

1. ✔ Dysplasia
2. ✘ Atypia
3. ✘ Anaplasia
4. ✘ Metaplasia

Question Number : 11 Question Id : 10922234342 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Pel Ebstein fever is characteristic of

Options :

1. ✘ Leukemia
2. ✔ Hodgkins lymphoma
3. ✘ Tuberculosis
4. ✘ Malaria

Question Number : 12 Question Id : 10922234343 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Causation of dental caries by *S.mutans* is facilitated by glucans due to

Options :

1. ✘ Glucans helps in the adhesion of the bacteria in the dental plaque
2. ✘ Glucans are resistant to hydrolytic enzymes, which causes plaque adherence
3. ✘ Glucans act as a barrier against the diffusion of the salivary buffers
4. ✔ All of the given options

Question Number : 13 Question Id : 10922234344 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Which of the following conditions is considered as cutaneous marker of breast cancer?

Options :

1. ✘ Plummer Vinson syndrome
2. ✘ Multiple endocrine neoplasia syndrome
3. ✔ Cowden syndrome
4. ✘ Bowen disease

Question Number : 14 Question Id : 10922234345 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Articular disc displacement is further classified by AAOFP as

Options :

- ✘ with reduction, without reduction
- ✘ without reduction, anteromedial or posteromedial displacement
- ✔ with reduction, without reduction, with intermittent locking
- ✘ anteromedial / anterolateral displacement, posteromedial / posterolateral displacement.

Question Number : 15 Question Id : 10922234346 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Hard tissue LASERS are useful for all the following purposes except

Options :

- ✘ Desensitise exposed root surfaces
- ✘ Make the hard tissue surfaces rough to promote bonding
- ✘ Vaporize the carious tissue
- ✔ To enhance pits and fissures

Question Number : 16 Question Id : 10922234347 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The correct sequence for multiple extractions include:

Options :

- ✔ Posterior teeth except 1st molar, anterior teeth except canine, 1st molar, canine
- ✘ 1st molar, posterior teeth, canine, anterior teeth
- ✘ Anterior teeth, canine, 1st molar, posterior teeth
- ✘ Canine, anterior teeth, posterior teeth, 1st molar

Question Number : 17 Question Id : 10922234348 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Dead space can be eliminated by the use of

Options :

- ✘ Single layer approximation
- ✔ Surgical packing of the defect
- ✘ Irrigation
- ✘ Use of sutures

Question Number : 18 Question Id : 10922234349 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The most difficult impaction among the following is

Options :

- ✘ Mesioangular
- ✔ Distoangular
- ✘ Vertical
- ✘ Transverse

Question Number : 19 Question Id : 10922234350 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Difficult extraction is anticipated in third molars with

Options :

- ✘ fused conical roots
- ✘ thin, slender roots
- ✔ less than 1/3 rd root completion
- ✘ short completely formed roots

Question Number : 20 Question Id : 10922234351 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Class IV of ASA physical status scale include patients with

Options :

1. ✘ mild-moderate systemic disease
2. ✘ Severe systemic disease with functional limitations
3. ✔ Severe systemic disease, that is a constant threat to life
4. ✘ Moribund patient, not expected to survive 24 hours, irrespective of treatment

Question Number : 21 Question Id : 10922234352 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Position for direct laryngoscopy

Options :

1. ✘ Supine position
2. ✘ Extended head position at atlanto-occipital level
3. ✔ Neck flexion at lower cervical spine with extension of head at atlanto-occipital level
4. ✘ Flexion of lower cervical spine

Question Number : 22 Question Id : 10922234353 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Pulse oximeter helps in measuring

Options :

1. ✘ Oxygen saturation of the patient
2. ✘ Heart rate
3. ✔ Oxygen saturation and Heart rate of the patient
4. ✘ None of the given options

Question Number : 23 Question Id : 10922234354 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Complete analgesia and amnesia with N₂O is known as

Options :

1. ✔ Light anesthesia
2. ✘ Moderate analgesia
3. ✘ Dissociative analgesia
4. ✘ Analgesic anesthesia

Question Number : 24 Question Id : 10922234355 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

State of indifference and immobilization, where patient becomes deeply sedated, partially /wholly amnesic but is capable of obeying commands is

Options :

1. ✘ Deep sedation
2. ✘ Conscious sedation
3. ✘ Analgesic anesthesia
4. ✔ Neuroleptanalgesia

Question Number : 25 Question Id : 10922234356 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Under general anesthesia, protective reflexes are

Options :

1. ✘ present
2. ✔ suppressed
3. ✘ not altered
4. ✘ Varies with the agents used

Question Number : 26 Question Id : 10922234357 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Complications during tracheal intubation include all the following except

Options :

1. ✘ Aspiration of stomach contents
2. ✘ Rupture of trachea and bronchus
3. ✔ Tracheal collapse
4. ✘ Airway obstruction

Question Number : 27 Question Id : 10922234358 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Management of wheezing during administration of GA include

Options :

1. ✘ Establishing the cause
2. ✘ Deepen the level of anaesthesia
3. ✘ IV salbutamol administration
4. ✔ All of the given options

Question Number : 28 Question Id : 10922234359 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

In 2 rescuer cardiopulmonary resuscitation , ratio of compressions and ventilation is maintained at

Options :

1. ✘ 15:1
2. ✔ 5:1
3. ✘ 15:2
4. ✘ 5:2

Question Number : 29 Question Id : 10922234360 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Articular surfaces of TMJ are covered by

Options :

1. ✘ Hyaline cartilage
2. ✔ Fibrocartilage
3. ✘ Elastic cartilage
4. ✘ Any of the above

Question Number : 30 Question Id : 10922234361 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Reduction of dislocation of TMJ is done

Options :

1. ✘ without anesthesia
2. ✘ under local anesthesia
3. ✘ under general anesthesia / sedation with muscle relaxants

4. ✓ Any of the above depending on the amount of muscle spasm

Question Number : 31 Question Id : 10922234362 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

During normal healing, the fracture line increases in width about 2 weeks after reduction of fracture due to?

Options :

1. ✗ Incomplete healing
2. ✗ Improper reduction
3. ✓ Resorption of fractured ends
4. ✗ Distraction

Question Number : 32 Question Id : 10922234363 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Recommended approach for condylar neck and ramus surgeries

Options :

1. ✓ Post ramal (Hind) approach
2. ✗ Endaural approach
3. ✗ Preauricular approach
4. ✗ Post auricular approach

Question Number : 33 Question Id : 10922234364 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The minimum recommended width of gap in gap arthroplasty to prevent reankylosis?

Options :

1. ✓ 1 cm
2. ✗ 2 cm
3. ✗ 0.5 cm
4. ✗ 2.5 cm

Question Number : 34 Question Id : 10922234365 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Starch iodine test is diagnostic test for

Options :

1. ✗ Sjogren's syndrome
2. ✗ Ramsay Hunt syndrome
3. ✗ Melkersson Rosenthal syndrome
4. ✓ Frey's syndrome

Question Number : 35 Question Id : 10922234366 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Treatment of trigger points in myofascial pain include

Options :

1. ✗ Ultrasound therapy
2. ✗ Injection of LA
3. ✗ TENS
4. ✓ All of the given options

Question Number : 36 Question Id : 10922234367 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

----- genioplasty is used to correct short vertical facial height with deep bites.

Options :

1. ✓ Lengthening
2. ✗ Augmentation
3. ✗ Reduction
4. ✗ Straightening

Question Number : 37 Question Id : 10922234368 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Glasgow coma scale rates

Options :

1. ✗ Motor response
2. ✗ verbal response
3. ✗ Eye examination and response
4. ✓ All of the given options

Question Number : 38 Question Id : 10922234369 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Battle's sign is

Options :

1. ✗ Bruises over the nasal bridge
2. ✗ Periorbital ecchymoses
3. ✓ Bruises behind the ear
4. ✗ Ecchymosis in floor of mouth

Question Number : 39 Question Id : 10922234370 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Biochemically, distinction between nasal secretion and CSF can be made as

Options :

1. ✗ CSF is rich in protein
2. ✓ CSF is rich in glucose
3. ✗ Nasal secretion is high in sugar
4. ✗ Nasal secretion is low in mucin

Question Number : 40 Question Id : 10922234371 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Absolute stability of fracture fragments is best achieved using

Options :

1. ✓ Open reduction
2. ✗ Maxillomandibular fixation
3. ✗ Intraoral traction
4. ✗ Extraoral traction

Question Number : 41 Question Id : 10922234372 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Circumferential wiring is done to treat edentulous jaws by the use of

Options :

1. ✗ Lateral compression splint
2. ✗ Stabilization splint
3. ✗ Repositioning splint

4. ✓ Gunning splint

Question Number : 42 Question Id : 10922234373 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Bone plate osteosynthesis is contraindicated in all the following except

Options :

1. ✗ Children with mixed dentition
2. ✗ Gross pathologies of the jaw
3. ✓ Edentulous patients with loss of bone segments
4. ✗ Badly comminuted fractures

Question Number : 43 Question Id : 10922234374 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Identify the wrong statement about compression plate system of osteosynthesis

Options :

1. ✗ Heals by primary healing of bone
2. ✓ The intrafragmentary gap should be less than equal to 0.8cm
3. ✗ Fragments are firmly approximated under pressure against each other
4. ✗ Precision is required in this technique.

Question Number : 44 Question Id : 10922234375 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Bones with least resistance to fracture in the middle third are

Options :

1. ✗ Zygomatic arch
2. ✓ Nasal bone
3. ✗ Maxilla
4. ✗ All of the given options

Question Number : 45 Question Id : 10922234376 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Upward displacement of fractured maxilla results in a

Options :

1. ✗ LeFort I fracture
2. ✗ Tripod fracture
3. ✗ LeFort IV fracture
4. ✓ Telescopic fracture

Question Number : 46 Question Id : 10922234377 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

LeFort IV fracture as per modification of LeFort's classification includes

Options :

1. ✗ LeFort I & II fractures and cranial base fracture
2. ✗ LeFort I & II fractures and nasal bone fracture
3. ✓ LeFort II & III fractures and cranial base fracture
4. ✗ LeFort II & III fractures and nasal bone fracture

Question Number : 47 Question Id : 10922234378 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Diplopia may be caused by

Options :

1. ✘ Muscle or periorbital entrapment in the orbital floor fracture
2. ✘ Displacement of the globe due to orbital floor fracture
3. ✘ Supranuclear impairments of 3rd, 4th, 6th cranial nerves
4. ✔ All of the given options

Question Number : 48 Question Id : 10922234379 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Image processing in digital imaging means

Options :

1. ✘ Any operation that converts a latent image to a visible image
2. ✔ Any operation that acts to improve, restore or analyse the image
3. ✘ Any operation that creates a latent image
4. ✘ All of the given options

Question Number : 49 Question Id : 10922234380 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Sharp instruments are best sterilized by

Options :

1. ✘ Autoclave
2. ✔ Hot air oven
3. ✘ Boiling water
4. ✘ All of the given options

Question Number : 50 Question Id : 10922234381 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Complete denture retention depends on

Options :

1. ✘ Extension of denture base
2. ✘ Intimate contact between the basal seat and denture base
3. ✘ Muscular factor
4. ✔ All of the given options

Question Number : 51 Question Id : 10922234382 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Sore spots seen in a edentulous patient given dentures 2 days back are due to

Options :

1. ✘ Unbalanced occlusion
2. ✘ Overextension of denture flanges
3. ✘ Mechanical irritation
4. ✔ All of the given options

Question Number : 52 Question Id : 10922234383 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Residual ridge reduction in mandible is

Options :

1. ✘ Greater than maxilla throughout
2. ✘ Lesser than maxilla throughout
3. ✔ Greater than maxilla in 1st year, lesser than maxilla in later years

4. ✘ Lesser than maxilla in 1st year, greater than maxilla in later years.

Question Number : 53 Question Id : 10922234384 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Residual ridge reduction is

Options :

1. ✔ Centripetal in maxilla, centrifugal in mandible
2. ✘ Centrifugal in maxilla, centripetal in mandible
3. ✘ Centripetal in both jaws
4. ✘ Centrifugal in both jaws

Question Number : 54 Question Id : 10922234385 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Chewing efficiency in edentulous patients may be enhanced by all the following except

Options :

1. ✘ Placement of implants
2. ✔ Preprosthetic surgical correction of ridges
3. ✘ Use of overdentures
4. ✘ Preserving a greater number of natural teeth

Question Number : 55 Question Id : 10922234386 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

In an edentulous patient, vertical dimension of occlusion and rest may be best determined by using

Options :

1. ✘ Making arbitrary marks on nose & chin
2. ✔ Anatomic reference points such as columella
3. ✘ Direct measurement between marked points
4. ✘ All of the given options

Question Number : 56 Question Id : 10922234387 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Deficiency in the amount of minerals relative to matrix in bone is

Options :

1. ✘ Osteoporosis
2. ✔ Osteomalacia
3. ✘ Osteosclerosis
4. ✘ Osteopetrosis

Question Number : 57 Question Id : 10922234388 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Retained roots should be left behind when they are

Options :

1. ✔ Difficult to distinguish from surrounding bone
2. ✘ Covered by mucosa only
3. ✘ Associated with cyst/ abscess
4. ✘ Surrounded by PDL

Question Number : 58 Question Id : 10922234389 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Surgical removal of tori is recommended in all the following conditions except

Options :

1. ✘ When they are sensitive to pressure
2. ✘ Extends beyond the vibrating line
3. ✘ Mandibular tori interfering with border seal
4. ✔ When located in midline on palate, irrespective of size

Question Number : 59 Question Id : 10922234390 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Parallelism of primary denture bearing areas affects

Options :

1. ✘ Denture retention
2. ✘ Denture esthetics
3. ✔ Denture stability
4. ✘ Denture adaptation

Question Number : 60 Question Id : 10922234391 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

An old male patient presented with a painless blue nodule of recent origin, with positive diascopy test, on the lower lip. The likely diagnosis is

Options :

1. ✘ Nevus
2. ✔ Varix
3. ✘ Vascular malformation
4. ✘ Amalgam tattoo

Question Number : 61 Question Id : 10922234392 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The function of surgical template in immediate dentures is

Options :

1. ✔ Guide to bone trimming
2. ✘ Guide to extraction
3. ✘ Guide for reduction of overdenture abutments
4. ✘ All of the given options

Question Number : 62 Question Id : 10922234393 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Overdentures are also known as all the following except

Options :

1. ✘ Teeth supported dentures
2. ✔ Non traditional dentures
3. ✘ Hybrid dentures
4. ✘ Assisted complete dentures

Question Number : 63 Question Id : 10922234394 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Midline fractures of dentures are caused predominantly due to

Options :

1. ✔ Flexural fatigue failure
2. ✘ Dropping of the denture

- ✘ Poor design of the denture
- ✘ Storing denture under dry conditions

Question Number : 64 Question Id : 10922234395 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The denture base material with incomplete polymerisation and inferior properties includes

Options :

- ✘ Heat activated resins
- ✔ Chemically activated resins
- ✘ Microwave activated resins
- ✘ Light activated resins

Question Number : 65 Question Id : 10922234396 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

All the following are true about porcelain teeth except

Options :

- ✘ Have better esthetic quality
- ✘ Are attached mechanically to resin base
- ✔ Can be reshaped to fit limited interarch spaces
- ✘ Cause significant wear of opposing natural teeth

Question Number : 66 Question Id : 10922234397 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Gelation of tissue conditioners is a

Options :

- ✔ Physical process
- ✘ Chemical process
- ✘ Physico-chemical process
- ✘ Biologic process

Question Number : 67 Question Id : 10922234398 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Advantages of base metal alloys as denture bases include

Options :

- ✘ Increased tongue space due to reduced bulk
- ✘ Increased weight to enhance stability
- ✘ Can withstand high masticatory stresses
- ✔ All of the given options

Question Number : 68 Question Id : 10922234399 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Vibrating line is located in

Options :

- ✘ Hard palate
- ✔ Soft palate
- ✘ Junction of hard and soft palate
- ✘ Buccopharyngeal raphe

Question Number : 69 Question Id : 10922234400 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Fracture of isthmus in an MO Amalgam filling is prevented by

Options :

1. ✘ Increasing the depth of cavity preparation
2. ✘ Unsupported enamel is removed at the cavosurface margin
3. ✘ If the proximal box is made larger and wider
4. ✔ Rounding of axiopulpal angle

Question Number : 70 Question Id : 10922234401 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The least retentive pins used in pin supported restorations is

Options :

1. ✘ Self threaded pins
2. ✔ Cemented pins
3. ✘ Friction locked pins
4. ✘ None of the given options

Question Number : 71 Question Id : 10922234402 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Corrosion that occurs between dissimilar metals in contact is

Options :

1. ✘ Electrochemical corrosion
2. ✔ Galvanic corrosion
3. ✘ Stress corrosion
4. ✘ Crevice corrosion

Question Number : 72 Question Id : 10922234403 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Deep caries is left over in which of the following techniques?

Options :

1. ✘ Always left behind
2. ✘ Direct pulp capping
3. ✔ Indirect pulp capping
4. ✘ Never left behind

Question Number : 73 Question Id : 10922234404 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The best way to protect pulp under any restoration is

Options :

1. ✘ Using liner
2. ✘ Using base
3. ✔ Intact dentin 2mm or more
4. ✘ All of the given options

Question Number : 74 Question Id : 10922234405 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Amalgam creep can be decreased by

Options :

1. ✘ High condensation forces
2. ✘ Accurate timing of trituration & condensation

- ✘ Use of high copper alloys
- ✔ All of the given options

Question Number : 75 Question Id : 10922234406 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Discoloured tooth is best conservatively treated by

Options :

- ✘ Crown
- ✔ Bleaching
- ✘ Veneer
- ✘ Extraction

Question Number : 76 Question Id : 10922234407 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

When there is an accidental mechanical pulp exposure, immediately

Options :

- ✔ The tooth should be isolated
- ✘ The tooth should be treated by endodontic therapy
- ✘ The tooth should be filled with amalgam
- ✘ None of the given options

Question Number : 77 Question Id : 10922234408 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The filling material with more microleakage is

Options :

- ✔ Composite
- ✘ Amalgam
- ✘ Direct gold filling
- ✘ Silicate cement

Question Number : 78 Question Id : 10922234409 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Reverse bevels in cast gold restorations are given to

Options :

- ✘ Improve aesthetics
- ✔ Counteract oblique forces
- ✘ Avoid high points
- ✘ All of the given options

Question Number : 79 Question Id : 10922234410 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

To record an impression after inlay preparation, in case of subgingival margins,----- is used

Options :

- ✘ Surgical exposure
- ✔ Gingival retraction cord with Alum, tannic acid
- ✘ Gingival retraction cord with ZOE
- ✘ All of the given options

Question Number : 80 Question Id : 10922234411 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Retention of cast gold restorations is not facilitated by

Options :

1. ✘ Cavo surface bevel
2. ✘ Near parallel walls
3. ✔ Undermining dentinal walls
4. ✘ Rounding of axiopulpal line angle

Question Number : 81 Question Id : 10922234412 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Cutting efficiency of bur is influenced by

Options :

1. ✘ Rake angle
2. ✘ Neck diameter
3. ✘ Spiral angle & cross cuts
4. ✔ All of the given options

Question Number : 82 Question Id : 10922234413 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Varnish is applied on only dentinal walls for

Options :

1. ✘ Composite
2. ✔ Cast gold restorations
3. ✘ Amalgam restorations
4. ✘ All of the given options

Question Number : 83 Question Id : 10922234414 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Primary resistance form in amalgam restorations is facilitated by all the following except

Options :

1. ✘ Box shape
2. ✘ Preservation of cusps and marginal angles
3. ✔ Use of pins
4. ✘ Rounding of internal line angles

Question Number : 84 Question Id : 10922234415 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Pulp vitality testing using blood supply of tooth include all the following except

Options :

1. ✔ Electric pulp testing
2. ✘ Pulse oximetry
3. ✘ Laser Doppler flowmetry
4. ✘ Thermographic Imaging

Question Number : 85 Question Id : 10922234416 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Endodontic instrument with maximum cutting efficiency is

Options :

1. ✘ K flex file
2. ✔ Hedstroem file

3. ✘ Unifile
4. ✘ Nickel-Titanium file

Question Number : 86 Question Id : 10922234417 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The file that does not undergo permanent deformation when used in curved canals is

Options :

1. ✔ Ni-Ti files
2. ✘ Stainless steel files
3. ✘ Ni-Ti files and Stainless steel files
4. ✘ None of the given options

Question Number : 87 Question Id : 10922234418 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Best technique to be used for canal preparation in fine, curved canals

Options :

1. ✘ Stepback method
2. ✔ Double flare technique
3. ✘ Hybrid technique
4. ✘ Stepdown technique

Question Number : 88 Question Id : 10922234419 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Single sitting RCT is indicated in all the following except

Options :

1. ✘ Tooth with pulpal exposure due to trauma
2. ✔ Teeth with infection
3. ✘ Teeth with internal resorption without external communication
4. ✘ All of the given options

Question Number : 89 Question Id : 10922234420 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Patient with an asymptomatic endodontically treated tooth 6 months back presents with a periapical radiolucency. Course of further action should be

Options :

1. ✘ Redo RCT
2. ✘ Perform periapical surgery
3. ✔ Observe
4. ✘ Extraction

Question Number : 90 Question Id : 10922234421 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Endodontic implants are used in

Options :

1. ✘ Periodontally involved teeth
2. ✘ Internal resorption affecting the strength of the root
3. ✘ Presence of 2 root fragments which cannot be aligned
4. ✔ All of the given options

Question Number : 91 Question Id : 10922234422 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Linear gingival erythema is seen in

Options :

1. ✘ Chronic desquamative gingivitis
2. ✔ HIV gingivitis
3. ✘ Plasma cell gingivitis
4. ✘ ANUG

Question Number : 92 Question Id : 10922234423 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Earliest clinical sign in gingivitis is

Options :

1. ✔ Bleeding on probing
2. ✘ Erthematous gingiva
3. ✘ Periodontal pocket formation
4. ✘ Reduction of stippling

Question Number : 93 Question Id : 10922234424 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Causes of gingival recession include all the following except

Options :

1. ✘ Faulty tooth brushing technique
2. ✘ Periodontal disease
3. ✘ Gingival ablation
4. ✔ Low frenal attachment

Question Number : 94 Question Id : 10922234425 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

In acute leukemia, gingival enlargement is due to

Options :

1. ✘ Deficiency of Vitamin C
2. ✔ Leukemic cellular infiltration
3. ✘ Altered response to plaque due to hormonal imbalance
4. ✘ Presence of local irritants

Question Number : 95 Question Id : 10922234426 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Punched out ulcers, papillary necrosis of gingiva, pseudomembrane formation in a 10 year old febrile patient is suggestive of

Options :

1. ✘ Candidiasis
2. ✘ Diphtheria
3. ✔ ANUG
4. ✘ Noma

Question Number : 96 Question Id : 10922234427 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Buttressing bone formation is seen in

Options :

1. ✘ Osteolytic lesions
2. ✔ Trauma from occlusion
3. ✘ Periodontitis
4. ✘ All of the given options

Question Number : 97 Question Id : 10922234428 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

DNA probes are used to

Options :

1. ✘ Calculate attachment loss
2. ✘ Calculate probing depth in cul de sac
3. ✔ Identify the presence of a specific microorganism
4. ✘ To assess the morphology of bone defects

Question Number : 98 Question Id : 10922234429 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Most effective and stable grasp for periodontal instrumentation is

Options :

1. ✔ Modified pen grasp
2. ✘ Palm and thumb grasp
3. ✘ Pen grasp
4. ✘ Finger grasp

Question Number : 99 Question Id : 10922234430 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Electrodes used for excision of tissue is

Options :

1. ✘ Loop electrodes
2. ✔ Single wire electrodes
3. ✘ Heavy electrode
4. ✘ All of the given options

Question Number : 100 Question Id : 10922234431 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Gingival sulcus is best cleaned by which of the following brushing technique?

Options :

1. ✘ Modified Stillman's method
2. ✘ Charters method
3. ✘ Fones method
4. ✔ Bass method

Question Number : 101 Question Id : 10922234432 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Gingivectomy is indicated in all the following conditions except

Options :

1. ✘ Removal of pseudopockets
2. ✘ Removal of gingival enlargements
3. ✔ Aesthetic regions
4. ✘ All of the given options

Question Number : 102 Question Id : 10922234433 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Long narrow defect on a single tooth is best treated by

Options :

1. ✘ Apically displaced flap
2. ✔ Laterally displaced flap
3. ✘ Coronally displaced flap
4. ✘ Papillary preservation flap

Question Number : 103 Question Id : 10922234434 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

With apically displaced flaps, the width of attached gingiva is increased by approximately ----- the pretreatment pocket depth

Options :

1. ✘ One third
2. ✘ Three- quarters
3. ✔ Half
4. ✘ The same as

Question Number : 104 Question Id : 10922234435 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Among the following autografts, the best one utilised for reconstruction is

Options :

1. ✔ Iliac crest graft
2. ✘ Costochondral graft
3. ✘ Metatarsal bone graft
4. ✘ Calvarial graft

Question Number : 105 Question Id : 10922234436 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The advantage of Biobrane is

Options :

1. ✘ Permits guided tissue regeneration
2. ✘ Biodegradable
3. ✘ Does not require removal after 4 weeks
4. ✔ All of the given options

Question Number : 106 Question Id : 10922234437 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Treatment advised for Grade II furcation involvement includes all the following except

Options :

1. ✔ Tunnelling
2. ✘ Root resection
3. ✘ GTR with bone grafts
4. ✘ Osteoplasty

Question Number : 107 Question Id : 10922234438 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Bone graft with osteoinductive property

Options :

1. ✘ Autogenous cancellous bone
2. ✔ Decalcified freeze-dried bone
3. ✘ Hydroxyapatite
4. ✘ Tricalcium phosphate

Question Number : 108 Question Id : 10922234439 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Which suturing technique is indicated when there is need for coverage of interdental bone by interdental papilla

Options :

1. ✘ Figure of eight
2. ✘ Anchor
3. ✘ Periosteal
4. ✔ Direct

Question Number : 109 Question Id : 10922234440 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Action taken to halt the progress of a disease at its incipient stage is

Options :

1. ✘ Tertiary prevention
2. ✔ Secondary prevention
3. ✘ Primary prevention
4. ✘ Primordial prevention

Question Number : 110 Question Id : 10922234441 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Manufactured preparation containing crushed betel nut, tobacco and flavouring agents is

Options :

1. ✔ Gutka
2. ✘ Khaini
3. ✘ Zarda
4. ✘ Mishri

Question Number : 111 Question Id : 10922234442 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

A tooth showing decay below a restoration is counted under ---- in DMFT index

Options :

1. ✔ D
2. ✘ M
3. ✘ F
4. ✘ Both D & F

Question Number : 112 Question Id : 10922234443 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Topical fluoride preparation that is unstable and needs to be prepared just before use is

Options :

1. ✘ 1.23% APF
2. ✘ 2% NaF

3. ✓ 8% SnF
4. ✗ All of the given options

Question Number : 113 Question Id : 10922234444 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Human anatomic waste should be disposed in ---- coloured bag

Options :

1. ✗ Red
2. ✗ Black
3. ✗ Blue
4. ✓ Yellow

Question Number : 114 Question Id : 10922234445 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Caries activity tests done by dental professional are a form of ---- prevention by a dental professional

Options :

1. ✓ Primary
2. ✗ Secondary
3. ✗ Tertiary
4. ✗ Quaternary

Question Number : 115 Question Id : 10922234446 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Twice daily brushing with fluoride paste and once without paste is recommended for children

Options :

1. ✗ Less than 4 yrs
2. ✓ 6-10 years
3. ✗ Above 10 years
4. ✗ 4-6 years

Question Number : 116 Question Id : 10922234447 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Rule of 10 in cleft lip repair includes

Options :

1. ✗ Hb of 10gm%
2. ✗ Weight 10 pounds, age 10 weeks
3. ✗ TLC less than 10,000/cu mm
4. ✓ All of the given options

Question Number : 117 Question Id : 10922234448 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

An uncooperative pediatric patient with multiple carious teeth may be best managed by

Options :

1. ✗ Conscious sedation
2. ✗ Premedication
3. ✓ General anesthesia
4. ✗ Reinforcement

Question Number : 118 Question Id : 10922234449 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Type of fear most difficult to overcome in children is

Options :

1. ✘ Innate fear
2. ✔ Subjective fear
3. ✘ Objective fear
4. ✘ All of the given options

Question Number : 119 Question Id : 10922234450 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

All the below are correct about HOME except

Options :

1. ✘ It is a kind of aversive conditioning
2. ✔ It is effective in children under 3 years
3. ✘ Indicated in children displaying uncontrollable behaviour
4. ✘ Indicated in children who can understand simple verbal commands

Question Number : 120 Question Id : 10922234451 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Mothers of which age group are allowed to stay in the dental operator

Options :

1. ✔ Under 2 years
2. ✘ Under 6 years
3. ✘ Under 10 years
4. ✘ Under 12 years

Question Number : 121 Question Id : 10922234452 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

In a child with rampant caries, all the following should be done in the first visit except

Options :

1. ✘ Caries excavation & restoration
2. ✘ Parent counselling
3. ✘ Topical fluoride application
4. ✔ Extraction of unrestorable teeth

Question Number : 122 Question Id : 10922234453 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Calcific bridge in a calcium hydroxide pulpotomy is formed from

Options :

1. ✘ Calcium hydroxide paste used
2. ✔ Blood vessel borne
3. ✘ Debrided cells
4. ✘ Any of the above

Question Number : 123 Question Id : 10922234454 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Following pulpectomy in a primary tooth, the canals are obturated with

Options :

1. ✘ Gutta percha
2. ✘ Calcium Hydroxide

3. ✓ ZOE

4. ✗ Formocresol

Question Number : 124 Question Id : 10922234455 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Non vital immature permanent tooth should be treated by

Options :

1. ✗ Indirect pulp capping

2. ✗ Calcium hydroxide pulpotomy

3. ✗ Apexogenesis

4. ✓ Apexification

Question Number : 125 Question Id : 10922234456 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Late mesial shift is accommodated by

Options :

1. ✗ Closure of primate space

2. ✓ Closure of leeway space

3. ✗ Closure of interdental spaces

4. ✗ All of the given options

Question Number : 126 Question Id : 10922234457 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Frenectomy in a case of midline diastema due to thick maxillary labial frenum evident by positive tension test should be done

Options :

1. ✗ Prior to midline diastema closure

2. ✗ After the closure of diastema

3. ✓ After the eruption of maxillary canines

4. ✗ Before the eruption of maxillary canines

Question Number : 127 Question Id : 10922234458 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Moyers analysis predicts

Options :

1. ✗ Width of all 4 mandibular incisors

2. ✗ Width of maxillary canine and premolars

3. ✗ Width of mandibular canine & premolar

4. ✓ Width of both maxillary & mandibular canines & premolars

Question Number : 128 Question Id : 10922234459 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

All the following procedures require no retention except

Options :

1. ✗ Anterior cross bite

2. ✗ Posterior crossbite

3. ✓ Severe rotation

4. ✗ Serial extraction

Question Number : 129 Question Id : 10922234460 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

After extraction of first premolars, the space will be closed down by movement of anterior & posterior teeth in ____ ratio

Options :

1. ✘ 40:60
2. ✔ 60:40
3. ✘ 30:70
4. ✘ 70:30

Question Number : 130 Question Id : 10922234461 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

The decreasing order of teeth in terms of resistance to orthodontic movement is

Options :

1. ✔ Mandibular molars > Maxillary cuspids > Mandibular cuspids > Maxillary molars
2. ✘ Maxillary Molars > Mandibular molars > Maxillary canines > mandibular canines
3. ✘ Max molar > Mand molar > Mand canine > Max canine
4. ✘ Max canine > Max molar > Mand molar > Mand canine

Question Number : 131 Question Id : 10922234462 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Myofunctional appliance used to treat maxillary deficiency is

Options :

1. ✘ Chin cap
2. ✔ Reverse pull headgear
3. ✘ Cervical pull headgear
4. ✘ Occipital pull headgear

Question Number : 132 Question Id : 10922234463 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

In stage II of Begg's technique, which of the following is done

Options :

1. ✘ Crowding of teeth is corrected
2. ✘ Inclination of teeth is corrected
3. ✔ Extraction spaces are closed
4. ✘ Anterior space closure

Question Number : 133 Question Id : 10922234464 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Pick up the true statement regarding orthodontic treatment

Options :

1. ✘ Greater the length of the wire, greater the force, faster the movement
2. ✘ Greater the diameter of the wire, greater the force, faster the movement
3. ✔ Greater the length of the wire, gentler the force, faster the movement
4. ✘ None of the given options

Question Number : 134 Question Id : 10922234465 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Even when GFR is less than 10ml/min, which of the following drugs may be given unchanged

Options :

1. ✘ Gentamicin
2. ✘ Ampicillin
3. ✔ Narcotics
4. ✘ Amphotericin

Question Number : 135 Question Id : 10922234466 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

When there is active bleeding in a case of undiagnosed bleeding disorder, the best treatment option is

Options :

1. ✘ Platelets
2. ✔ Fresh frozen plasma
3. ✘ Cryoprecipitate
4. ✘ Factor concentrates

Question Number : 136 Question Id : 10922234467 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

All the following should be followed for a hyperthyroid patient except

Options :

1. ✔ Surgery need not be discarded as there is no bleeding risk
2. ✘ Epinephrine is contraindicated
3. ✘ Short time appointments are recommended
4. ✘ Propylthiouracil may cause sialolith formation

Question Number : 137 Question Id : 10922234468 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

In patients receiving long term steroid therapy, if invasive surgery is planned

Options :

1. ✔ 100-200mg hydrocortisone should be administered to prevent adrenal crisis on the day of surgery
2. ✘ No additional steroid cover is recommended
3. ✘ Daily dose of steroids should be stopped prior to surgery
4. ✘ No additional steroid cover is required if daily dose is less than 40mg prednisone

Question Number : 138 Question Id : 10922234469 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Adynamic bone disease is characterised by

Options :

1. ✘ Reduced osteoblasts and osteoclasts
2. ✘ Low burn turnover
3. ✘ Low levels of PTH
4. ✔ All of the given options

Question Number : 139 Question Id : 10922234470 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Imaging technique that estimates internal bone density in Hounsfield units is

Options :

1. ✘ CBCT
2. ✔ CT
3. ✘ OPG
4. ✘ All of the given options

Question Number : 140 Question Id : 10922234471 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

During implant planning, the distance between 2 implants should be kept at a minimum of

Options :

1. ✘ 1.5 mm
2. ✘ 2mm
3. ✘ 2.5mm
4. ✔ 3mm

Question Number : 141 Question Id : 10922234472 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

First radiographic sign of osteomyelitis is

Options :

1. ✘ Multiple irregular radiolucent areas
2. ✘ Onion peel appearance
3. ✔ Blurred or fuzzy trabeculae
4. ✘ Erosion of cortex

Question Number : 142 Question Id : 10922234473 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Which of the following conditions produces snowcapped appearance of teeth

Options :

1. ✔ Amelogenesis imperfecta
2. ✘ Dentinogenesis imperfecta
3. ✘ Dentin dysplasia
4. ✘ Regional Odontodysplasia

Question Number : 143 Question Id : 10922234474 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Pumping tooth syndrome with soap bubble radiolucency and sunburst appearance is seen in

Options :

1. ✘ Osteosarcoma
2. ✔ Central hemangioma
3. ✘ Neurofibromatosis
4. ✘ Intraosseous carcinoma

Question Number : 144 Question Id : 10922234475 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Trap door is a radiological sign of

Options :

1. ✘ Le Fort I
2. ✘ Le Fort II
3. ✔ Orbital blow out
4. ✘ Le Fort III

Question Number : 145 Question Id : 10922234476 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Tree in winter and bush in winter are sialographic appearances of

Options :

1. ✘ Sjogren's syndrome
2. ✔ Normal parotid & submandibular glands
3. ✘ Salivary gland tumor
4. ✘ Sialadenitis

Question Number : 146 Question Id : 10922234477 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Digital autopsy facilitates

Options :

1. ✘ 3 D visualisation of a bullet path
2. ✘ Non destructive documentation process
3. ✘ Visualisation of soft tissue and bone
4. ✔ All of the given options

Question Number : 147 Question Id : 10922234478 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Bulls eye or target appearance is seen in

Options :

1. ✘ Erythema multiforme
2. ✘ Buccolingual dilacerations
3. ✘ Phleboliths
4. ✔ All of the given options

Question Number : 148 Question Id : 10922234479 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Bright signal on T2 weighted image seen in joint space is suggestive of

Options :

1. ✘ Osteoarthritis
2. ✔ Effusion
3. ✘ Ankylosis
4. ✘ Internal derangement

Question Number : 149 Question Id : 10922234480 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

HSV can be differentiated from VZV by all the following except

Options :

1. ✘ PCR
2. ✔ Cytology
3. ✘ Direct Fluorescent antigen detection
4. ✘ Antibodies

Question Number : 150 Question Id : 10922234481 Display Question Number : Yes Single Line Question Option : No Option Orientation : Vertical

Radiological signs of fracture include all of the following except

Options :

1. ✔ One or two sharply defined radiolucent lines extending beyond the anatomic boundary of a structure
2. ✘ Change in the normal anatomic outline or shape of the structure
3. ✘ Loss of continuity of outer border
4. ✘ Increase in radiopacity of a structure